

Yale University

## EliScholar – A Digital Platform for Scholarly Publishing at Yale

---

Yale Medicine Thesis Digital Library

School of Medicine

---

11-3-2006

# The Effects of Cyclic Strain on Rat Tail Tenocytes

Richard James Crockett  
*Yale University*

Follow this and additional works at: <http://elischolar.library.yale.edu/ymtdl>

---

### Recommended Citation

Crockett, Richard James, "The Effects of Cyclic Strain on Rat Tail Tenocytes" (2006). *Yale Medicine Thesis Digital Library*. 232.  
<http://elischolar.library.yale.edu/ymtdl/232>

This Open Access Thesis is brought to you for free and open access by the School of Medicine at EliScholar – A Digital Platform for Scholarly Publishing at Yale. It has been accepted for inclusion in Yale Medicine Thesis Digital Library by an authorized administrator of EliScholar – A Digital Platform for Scholarly Publishing at Yale. For more information, please contact [elischolar@yale.edu](mailto:elischolar@yale.edu).

The Effects of Cyclic Strain on Rat Tail Tenocytes

A Thesis Submitted to the  
Yale University School of Medicine  
In Partial Fulfillment of the Requirements for the  
Degree of Doctor of Medicine

by

Richard James Crockett

2006

## **Abstract**

### THE EFFECTS OF CYCLIC STRAIN ON RAT TAIL TENOCYTES

Richard Crockett, Michael Centrella, Thomas McCarthy, J. Grant Thomson. Section of Plastic Surgery, Department of Surgery, Yale University, School of Medicine, New Haven, CT.

The purpose of this study was to examine the effects of cyclic tension on the expression of hyaluronic acid, its receptor (CD44), and total glycosaminoglycan content in tendon fibroblasts. An *in vitro* model was used to analyze tenocytes from the tail tendons of rats. Tenocytes in the experimental group were exposed to cyclic mechanical stress, and using ELISA, western blot, and colorimetric dye-binding assays, the effect of strain on cultured tenocytes was examined.

Tenocytes exposed to mechanical strain produced  $1528 \pm 58$  ng/mL (mean  $\pm$  SEM) of hyaluronic acid, while those in a control environment produced only  $730 \pm 27$  ng/mL; nearly a two-fold difference ( $p < .0001$ ,  $n=44$ ). CD44, the receptor for hyaluronic acid, was also detected in higher concentrations. Tenocytes under mechanical strain increased their concentration of CD44 by 62.5%, with tenocytes exposed to strain having an optical density of  $26 \cdot 10^3 \pm 2 \cdot 10^3$  compared with  $15 \cdot 10^3 \pm 1 \cdot 10^3$  in controls ( $p < .05$ ,  $n=6$ ). The total glycosaminoglycan content of the two groups did not differ significantly; strained cells produced  $10.2 \pm 0.6$   $\mu$ g/mL and controls producing  $15.3 \pm 3$   $\mu$ g/mL ( $p=0.103$ ,  $n=44$ ).

We conclude that mechanical strain in tendon fibroblasts is sufficient to induce the production of hyaluronic acid and increase the expression of its receptor, CD44. The results of our study suggest that the beneficial effects on tendon adhesion formation seen with both mechanical strain and hyaluronic acid may be related in their mechanism.

## **Table of Contents**

<i>Introduction</i> .....	1
<i>Materials and Methods</i> .....	18
<i>Results</i> .....	25
<i>Discussion</i> .....	28
<i>References</i> .....	33

## **Acknowledgements**

I would like to thank my thesis advisor, J. Grant Thomson, MD, for his continued support, advice, and commitment during the course of this project. In the laboratory, Thomas McCarthy, PhD and Michael Centrella, PhD were an always-available and invaluable resource as I navigated the world of bench science.

Also, I want to express my gratitude to Martijn van Griensven, MD, PhD, Tanja Barkhausen, PhD, and Michael Skutek, MD, from Hannover, Germany who took the time to teach me the intricacies of tendon fibroblast tissue culture.

Finally, this work would not have been possible without the financial support of the Yale University Section of Plastic Surgery and the Office of Student Research. Thank you for investing in what I hope is a valuable contribution to our understanding of tendon biology.

## **Introduction**

No subject has created more challenges, interest, or discussion in the field of hand surgery than flexor tendon injuries. Controversies on the proper approach to tendon surgery began as early as Galen and Avicenna, and by the 1930s, Bunnell had coined the term “No Man’s Land” to describe a portion of the hand in which tendon injuries caused particular tribulation.(1)

The great difficulty with repairing flexor tendons surgically is the frequent creation of fibrotic adhesions between the tendon and surrounding tissues. These accumulations of scar tissue prevent the smooth gliding of the tendons, limiting movement and causing debilitating contractures that are the bane of hand surgeons.

The challenge presented by adhesions has been the focus of many research endeavors. Unfortunately, to date these have met with limited success. Are they truly a “biologic inevitability” as some authors(2) have described them?

Adhesions have long been an obstacle to proper healing after tendon injury or repair(3) and have many causes.(4) They differ in their degree of density, and different types of adhesions may be caused by independent regulatory pathways. This has led to attempts by some researchers to develop a standard methodology of measuring adhesion biomechanical strength.(5) Dense adhesions, for example, are characterized not only by an increased quantity of collagen than loose adhesions, but also with an increase in the ratio of type III to type I collagen and their ratio of cross-linking.(6) In tendons,

fibroblasts from the fibro-osseous sheath have biochemical and cellular differences which may enable them to migrate and contribute to their role in adhesion formation.(7)

Many techniques have been developed to mitigate the development of adhesions in tendons including: enveloping the repaired tendons within silicone sheathes(8), fibrin glues(9, 10), iontophoresis(11), pulsed electromagnetic fields(12, 13), and thermal preconditioning to induce the formation of cytoprotective heat shock proteins(14, 15). The results have not translated into successful clinical applications.

A number of biochemical agents have also been used to thwart the formation of adhesions. Non-steroidal anti-inflammatory agents that inhibit cyclooxygenase (COX) and in turn, prostaglandin synthesis, have received some attention. Indomethacin has been shown to be beneficial at reducing adhesions(16-19), and some work shows that ibuprofen can also reduce adhesions in both oral(20) and injectable(21) form.

Conversely, prostaglandins are known to cause or facilitate adhesion formation.(22) Corticosteroids(23, 24) and promethazine(25, 26) have also been seen to exert a favorable effect. Some of the benefits to pharmacological anti-inflammatory therapy may be due to the decreased edema in tissues. This is supported by biomechanical animal studies which examine the time course of gliding function after injury and reveal a significant impediment to gliding within hours of injury, long before adhesion formation can occur.(27)

Intraoperative exposure of repaired tendons to 5-fluorouracil has been shown to diminish adhesion formation in rabbit(28, 29) and chicken(30) models, as well as reducing the level of TGF $\beta$ 1 (thought to be involved in the formation of adhesions).(29) Later research has shown that this does not decrease the strength of the tendon repair.(31)

This may be due to its inhibitory effects on matrix metalloproteinases (particularly mmp2 and 9).(32) Aprotinin, another protease inhibitor, has similarly been shown to be effective in reducing adhesion formation(33) as has halofuginone, an inhibitor of collagen I transcription(34) and Beta-aminopropionitrile (BAPN), an inhibitor of collagen maturation.(35) The effect of these chemicals is to hinder collagen production, a needed component for the creation of adhesions. Collagen itself, when added topically to injured tendons, induces the upregulation of collagenase to mediate these effects.(36-38)

Although some of these have shown promise, the data are usually limited to small case series or animal models with clinical trials often yielding disappointing results. Despite some indications to the contrary(39, 40), physical barriers to adhesion formation such as PTFE have also not proven a useful technique(41-45). The use of an anti-adhesive membrane (ADCON-T/N) was evaluated in human flexor tendon repair in a prospective double-blind randomized control trial (RCT), and total active motion was evaluated at 3, 6, and 12 months after the repair. Although the length of time taken to achieve the final range of motion was shorter in treated patients, the total range of motion achieved was not improved.(46) The failure of ADCON-T/N to prevent adhesions has been supported by other researchers.(47)

As a whole, these studies emphasize that there are a variety of factors which may influence the formation of adhesions after tendon surgery. Along with these biochemical signals, mechanical strain also seems to be important for the proper functioning and healing of tendons. The tension-bearing property of tendons is important for the normal mechanics of joint motion. Unfortunately, after surgery tendon motion is limited by pain and rupture rates can be high if motion is allowed. Depending on the particular repair

technique and mobilization protocol, the rupture rates of digital flexor tendons range from 4%(48) to 46%(49) and 17% for thumbs(50). On average, the rupture rate approaches 10%(48, 50-65), and the rate of “excellent” or “good” results (using Strickland’s revised scoring technique, a commonly used tool to evaluate outcomes of hand surgery(53)) is often below 75%, although individual results vary considerably.

Historically, hand immobilization was used to protect flexor tendons after repair. This period often lasted three or more weeks, and while effective at minimizing the risk of tendon rupture, it had the disadvantage of maximizing the formation of adhesions. More recently, a substantial body of literature describing the beneficial role of early hand mobilization has been published. We now know that mobilization is beneficial to the smooth gliding of tendons and prevention of adhesions. And although active mobilization can result in rupture(66) or gap formation(67) in animal models, most data suggest a positive role. Overall, animal models using active rehabilitation mobilization protocols have been shown to decrease adhesion formation without undue risk of rupture when compared to immobilization.(68) This translates into repairs which are stronger, have a more efficient healing process(69) and greater range of motion(70) in the joints that they affect.

The timing of when tendon mobilization is started after surgery is also an important factor. In a study that examined canine flexor tendons after surgical repair, immediate mobilization was superior to delayed mobilization in returning range of motion (a 50% increase) and dramatically better than immobilized tendons (more than a five-fold improvement in angular rotation).(71) This benefit has been corroborated by other studies.(72)

There is still some discussion on which protocols maximize the beneficial effects of mechanical strain while introducing the least risk of rupture. It is known that several parameters play a role in determining outcome. The frequency of motion, for example, is an important factor, with increased frequency correlated with improved tensile strength.(73) A prospective multi-center RCT was performed to examine the effects of daily passive motion and whether increased duration of motion could improve tendon gliding after flexor tendon repair. The authors report that there was a significant benefit to increased duration of controlled motion on the range of motion, increasing the angle of rotation by 19 degrees.(74)

This has also translated into clinical research(74-76), and it is now clinically accepted practice to apply these protocols to patient care as the most effective method of preventing adhesions(4) even in the presence of concurrent digital nerve injuries.(77)

Previously, it was thought that active mobilization protocols (involving the contraction of muscles in continuity with the repaired tendons) would lead to unacceptable rupture rates. Mobilization however, both active and passive, helps tendon healing and decreases adhesion formation. Animal studies have supported a benefit to tendon strength and smooth gliding by using controlled passive movement(78) as well as active movement. Prospective trials in humans have shown an advantage in patients who have passive(79) and active(57-59) post-operative mobilization. Although some authors report an advantage to active motion(62), at least some data suggest that neither is superior to the other.(49) Active motion can result in higher rupture rates, but an analysis of different protocols shows that the risk can be comparable to passive motion if

performed judiciously(63) and if the patient avoids imprudent acts which account for half of all tendon ruptures.(50)

Cyclic mechanical loading of tendons to mimic strains encountered during mobilization protocols reveals that tenorrhaphy failure rates may depend on the repair technique used.(80) Indeed, there is an entire body of literature which examines the surgical techniques of tendon repair(42) which will not be discussed at length here except to say that after tendon injury or repair, there is an increase in the friction of tendon gliding which remodels with time.(81) High friction suture techniques(82) and many other factors(83) may centrally contribute to adhesion formation and possibly rupture.

Fibroblasts show a response to mechanical strain *in vitro* at both a cellular and biochemical level.(84) Tendons *in vivo* exposed to extracorporeal shock waves after injury have been shown to have an increased number of capillaries and decreased formation of adhesions, with a correspondingly higher force required to cause rupture.(85)

Cyclical mechanical strain, in particular, seems to be important. When exposed to cyclic mechanical strain, flexor tendons fibroblasts proliferate and align in the direction of tension to form a thicker growth layer.(86) Constant strain, however, appears to be of lesser benefit in healing tendons.(87, 88)

One question that exists regarding the effect of mechanical strain on healing tendons is the degree to which strain actually contributes to the improvements seen with active and passive motion rehabilitation protocols. One theory might argue that only motion is important, as the primary mechanism is the mechanical separation of fibrotic

bridges which are putative adhesions. Another view is that the strain is primarily important, as the host of growth factors regulated by and affecting strain certainly plays a role.

The question was addressed by a study which examined *in vivo* chicken flexor tendons. Motion and strain each had an independent roll in generating tensile strength, and the combination of the two was greater than either alone. An increase in collagen production, however, was seen only when tendons experienced some type of strain.(89) Other studies have shown that the degree of excursion (change in joint angle) is of minimal importance in determining the extent of adhesion formation or whether a repair will have good range of motion.(90)

The effect of mechanical strain on the growth factor milieu of tendons is paramount. After tendon injury or repair, a host of growth factors are upregulated. Microarray analysis of rat flexor tendons under strain demonstrate an “antifibrotic” pattern of gene regulation, with decreased transcription of collagen I and III, FGF, PDGF, and IGF1.(91) IGF has been shown to promote fibroblast growth in culture.(92-94) In contrast, TGF $\beta$ 1 (which is often associated with increased adhesion formation) was found to be upregulated during strain in this study.(91) The import of this is somewhat unclear.

The regulation of the TGF $\beta$  pathway seems to be particularly important in the pathogenesis of fibrosis and formation of adhesions. It has been shown to be associated with adhesion formation, and some studies show that the addition of TGF $\beta$  to tenocytes *in vitro* can increase collagen production(95), while neutralizing antibodies to TGF $\beta$  can decrease collagen production(96). The role of individual isoforms of TGF $\beta$  (TGFB1, 2, and 3) are not yet clear. Most evidence points to TGF $\beta$ 1 acting to contribute to adhesion

formation, and less so for TGF $\beta$ 2 and 3. This has been supported by research showing decreased adhesion formation in the presence of TGF $\beta$ 1 blocking antibodies(97) and the observation that although TGF $\beta$ 1 is expressed in healthy tenocytes, the levels rise sharply after tendon injury.(98)

Some research suggests that the site of action of individual growth factors varies significantly. In one study which examined the effect of a tendon injury on growth factor regulation at ten days after injury, TGF $\beta$  was detected around the repair site and just proximal to it, while other factors such as VEGF and PDGF were expressed throughout the tendon. Some growth factors such as EGF, IGF, and bFGF were not seen in tenocytes; only in pro-inflammatory cells in the repair site.(99) This increase in TGF $\beta$  activation is most pronounced in the tendon sheath and epitenon, peaking within two weeks of injury or repair.(100)

The expression of VEGF in injured tendons suggests that the supply of nutrients and oxygen may be an important factor in determining the adhesion forming response of healing tendons. When tendons are examined after repair, progressive changes in the microvasculature can be observed.(101) Revascularization of tendons following injury occurs progressively over a 7-21 day window, forming an extrinsic vascular supply.(102) In one study, the authors noted a decreasing density and more longitudinal orientation of vessels associated with an increasing range of motion after the acute healing phase of the first 21 days had passed.(103)

Other vascular signaling intermediaries have been implicated as well. Nitric oxide synthase (NOS) levels double after an acute tendon injury, and appear to be integral to the repair process. Blocking NOS production creates a persistent, chronic

inflammatory response that leads to an increased incidence and severity of adhesions.(104) Lactate can also modulate early wound healing and can stimulate collagen production after injury. Cultured tendon cells exposed to lactate have been shown to increase collagen production(105), suggesting that lactate levels may directly modulate collagen production during tendon wound healing or repair.

As newer molecular biology techniques become available, they will advance our understanding of the complex growth factor regulation that plays a critical role in the biology of adhesion formation and tendon healing. Adenovirus mediated gene transfer appears to be an efficient, dose dependent method of delivering transgenes to flexor tendons and at viral concentrations of less than  $10^9$ , does not appear to engender a significant inflammatory response.(106) Also, antisense oligonucleotides to procollagen have been shown effective at reducing collagen production in human tenocytes.(107)

In sum, many applications for the prevention of adhesions after tendon repair have been tried. Unfortunately, few have met with clinical success. To understand the biological basis of adhesions, a closer look at tendon biology is warranted.

Tendon is specialized connective tissue specifically designed to withstand tension and allow the smooth gliding of tissues. On gross examination, tendons themselves are bright white cord or band-like structures that form the structural connections between bone and muscle. Water constitutes approximately 70% of tendon mass(108) with collagen I accounting for the majority of the dry mass. Tendons contain additional structural elements, including bone attachment and insertion sites, blood vessels, and a dense extracellular matrix. The primary role of tendons is structural, which is reflected in

their decreased metabolic activity. Oxygen consumption, for example, is 7.5 fold lower when compared with skeletal muscles.(109) This property allows them to function in relatively ischemic environments; an important adaptation as tendons are commonly in compact anatomical regions with limited tissue perfusion and often under enormous tension which would otherwise cause tissue necrosis. However, the decreased metabolic rates also contribute to slower healing after a traumatic insult.(110)

Although tendons contain some elastin, they are comprised predominantly of thick collagen type I fibers and spindle-shaped tenocytes interspersed in a columnar fashion among the fibers. The glycosaminoglycan hyaluronic acid (hyaluronan, hyaluronate) is present in small quantities. External to the collagen fibers is a layer of paratenon fibroblasts. The internal fibroblast may be responsible for the bulk of the collagen synthesis, whereas the external cells may provide lubrication for tendon gliding during motion.(111)

At a micro-structural level, tendons are formed by the successive grouping of larger and larger bundles of fibers. At the smallest level, a triple-helix polypeptide chain forms “primary bundles” of tropocollagen which are only 15 angstroms in diameter. Several small fibers of tropocollagen together form a 35 angstrom microfibril, many of which unite to form a subfibril of 100-200 angstroms in width. Subfibril groups form 500-5000 angstrom fibrils, and eventually the 50-300  $\mu\text{m}$  fascicles of tendons.(112)

Each bundle of tendon collagen is covered by an *endotenon*. The septa of these endotenon form together to constitute an external *epitenon* that covers the three-dimensional surface of the tendon. This is further complemented in the hand by a thin adventitia known as the *paratenon* within a synovial-like fluid environment.(42)

This arrangement is coupled by a complex waveform or crimp structure of different topographies that forms a dynamic rope-like tendon capable of tremendous tensile strength. Biomechanical analyses, for example, reveal that tendons can withstand forces equivalent to thousands of Newtons.(113) The capacity to produce such a robust, complex structure lies in the equally intricate arrangement of cellular subtypes.

Tendons contain several types of cells. Tenocytes (or tendon fibroblasts) account for 90 to 95% of tendon cells. Synovial cells, chondrocytes located near osseous insertions sites, endothelial cells, smooth muscle cells of arterioles, and immunologic cells form the remainder. Collectively, they are responsible for the formation, maintenance and interactions of the extracellular matrix.

Along with the previously mentioned cellular components, extracellular fluid, and collagen and elastin fibers, the extracellular matrix (ECM) contains a fourth component – the ground substance. Ground substance is the portion of the extracellular matrix comprised of glycoproteins and proteoglycans – large molecules which are formed by the union of a glycosaminoglycan through a trisaccharide link to a protein core. Glycosaminoglycans (or GAGs) are synonymous with “mucopolysaccharides” and exist independently in the extracellular matrix as the most plentiful heteropolysaccharides in the body. GAGs are formed as long polymers of modified disaccharides and uronic acid. Because of their large size (molecular weights in the tens of millions) and predilection for absorbing water, glycosaminoglycans often play a structural role in lubrication or minimizing mechanical compression.

One glycosaminoglycan of particular interest is hyaluronic acid (HA).

Hyaluronic acid is a high molecular weight glycosaminoglycan first described in 1934.(114) It is formed by the repeating disaccharide N-acetylglucosamine and D-glucuronate.(115) Because of its high molecular weight, random coiled structure, and ability to attract water, hyaluronic acid is hydrodynamic and forms solutions that have high viscosity, elasticity, and propensity for filtration.(114)

Hyaluronic acid is unique in that it is not sulfated and not covalently bonded to a protein, though it does complex with other proteoglycans. It lubricates tendon tissues and is thought to play an important role in the early stages of connective tissue healing and scarless fetal wound healing.(116-119) In an experimental model involving fetal lambs, tendons which were injured healed with no subcutaneous scarring, no adhesions, and regained a smooth gliding surface with well organized collagen architecture, although adult sheep controls experienced significant dense scarring and adhesions.(120)

Interestingly, it has been shown that acellular amnion membrane (which contains elevated concentrations of hyaluronic acid(121)) can prevent the adhesion of tendons after repair without compromising tendon healing.(122, 123) In one study, the combined effect of amniotic membranes applied in conjunction with hyaluronic acid was examined. Although the researchers reported that this was greater than either amniotic membrane or hyaluronic acid alone, it did not reach statistical significance.(124) Other studies, however, have shown in animal models that topical amniotic fluid alone applied during tendon repair is sufficient to both decrease adhesion formation and improve the tensile strength of the repair.(125)

In animal models of osteoarthritis, the addition of hyaluronic acid has been shown to provide a lubricating effect, decreasing the friction of tissues.(126) Even in healthy joints, the lubricating effect of hyaluronic acid can be seen.(127)

It has been suggested that the molecular weight of hyaluronic acid plays an important role in determining its effect on adhesion formation.(128) Some investigators have attempted to examine whether the molecular weight affects the mechanical friction in joints. In one study, the researchers injected 1% hyaluronic acid of two different molecular weights into the joint spaces of rabbits and determined that molecular weight did not play a role in determining the lubricating characteristics.(129) Unfortunately, the molecular weights chosen by the researchers were both similar (one and two million Daltons) and relatively low. As a consequence, they may not have been sufficiently different to allow for detection of lubricating or signaling potential. Moreover, it may not be the lubricating role of hyaluronic acid that is central to its effects on tendon healing.

Hyaluronic acid was previously thought to be an inert molecule that provided mechanical compression support in connective tissues. It is now understood that hyaluronan is an active ligand that exerts important regulatory and healing effects through hyaluronan-binding proteins, also known as hyaladherins.(114) These interactions make hyaluronic acid an important player in regulating cell mobility, adhesion, and proliferation; all important cellular functions in tendons. Hyaluronic acid is also present intracellularly and may be incorporated from extracellular stores(130), or produced *de novo* from within the cell(131) where it is thought to influence cell proliferation and motility.(114)

CD44, the primary receptor for hyaluronic acid(132), was first cloned in 1989.(133) Antibody mediated activation of the CD44 receptor produces an anti-inflammatory response(134) and organisms which are engineered to eliminate CD44 gene expression (-/-) are unable to properly respond to injury.(114) In addition, hyaluronic acid and its primary receptor, CD44, have been implicated as important cell adhesion mediators in the pathogenesis of cancer(135).

Of the many roles hyaluronic acid is thought to have, the most germane to tendons is its role as a structural lubricant and metabolite in the regulation of cell proliferation and adhesion formation. Applied topically during flexor tendon repair in animal models, hyaluronic acid has been shown to decrease adhesions at the repair site and synovial sheath. In a study involving the flexor tendons in dogs, for example, hyaluronic acid was applied topically to the second and fifth flexor tendons which had been transected and repaired. After five weeks of immobilization, the tendons exposed to hyaluronic acid had a significant decrease in both gross and histological adhesions.(136) This has also been seen in other studies of rabbits.(137)

After flexor tendon repair, canine tendons exposed to a hyaluronic acid-containing solution showed reduced excursion resistance.(138, 139) By decreasing the friction and force on the repaired tendon, increased gliding occurs, which decreases adhesion formation(140) and the likelihood of rupture.

Multiple *in vitro* and *in vivo* animal studies have supported a role for hyaluronic acid at preventing adhesions.(141, 142) Collagen contents (adhesions) in peri-tendon tissues are decreased, smooth gliding is improved(143), and range of motion is greater.(144) Hyaluronic acid has been shown to increase expression of VEGF and

collagen IV when applied to healing tendons after repair, both potential anti-adhesive mediators.(140)

Although some studies have not shown hyaluronic acid to decrease adhesion formation(145-147), they are in the minority. During tendon repair in a rabbit model, Seprafilm, a bioresorbable membrane containing hyaluronic acid and carboxymethyl cellulose, worked as well as hyaluronic acid gel in the prevention of adhesions and improvement of range of motion when compared to controls who received no hyaluronic acid.(148) Benefits are also observed for tenolysis of existing adhesions in a chicken model(149) and in dogs.(136) Other preparations of hyaluronic acid such as Healon and viscoelastic gel have been shown to be effective at reducing adhesion formation, even when injected into zone II flexor tendon repairs in immobilized primates.(150) The addition of a phospholipid lubricant, dipalmitoyl phosphatidylcholine, has also been shown to augment the action of hyaluronic acid, decreasing the friction coefficient and formation of adhesions in rabbit flexor tendons.(151) Exogenously applied hyaluronic acid has also been shown to inhibit tendon fibroblast proliferation involved in the formation of adhesions in rabbit models.(152)

A potential role for hyaluronic acid to mitigate adhesion formation is of particular interest in hand surgery as adhesions can cause significant functional impairment after surgical tendon repair. Hyaluronic acid, although effective at reducing adhesions experimentally, has been disappointing clinically.(153, 154)

One of the challenges of applying hyaluronic acid as a therapy is its rapid absorption in the body. Hyaluronic acid can be degraded within a matter of hours, and within 24 hours, significant dilution may take place. Higher molecular weight

preparations and those that contain a higher initial concentration of hyaluronic acid experience a slower decline.(155, 156) This is further supported by evidence showing that hyaluronic acid which has been modified to persist longer has a beneficial effect. One group of researchers using a chicken model mixed hyaluronate with carboxymethylcellulose and prevented the formation of dense adhesions.(157)

A rabbit study which reported a beneficial effect from hyaluronic acid, decreasing adhesion formation from 95% to 45%, also noted that when adhesions did form in the presence of hyaluronic acid, they were less severe.(158) The authors show in a later study that improved function may have been a result of increased viscosity of the hyaluronic acid preparation.(159) Using *ex vivo* canine tendons as a model, a derivatized form of hyaluronic acid, which through chemical modification persists longer than native hyaluronate, maintained a lower gliding resistance over many repeated cycles of tendon movement than hyaluronic acid or saline.(82)

Clinically, some researchers have reported beneficial results with hyaluronic acid(160), but the vast majority have not.(42) This disparity between the very promising *in vitro* and animal study findings, and the disappointing clinical observations, is an area of great interest. A better understanding of the biology and regulation of hyaluronic acid is needed.

It has been shown that removing the tensional loading from tendons either experimentally or surgically changes matrix characteristics including increases in total glycosaminoglycan content.(111) Conversely, adding motion to repaired tendons has been shown to be beneficial to tendon healing, resulting in fewer adhesions, increased strength of repair, and more longitudinal alignment of collagen bundles compared with

healing tendons without movement.(54, 71) How tendon cells perceive changes in tensional loading, however, is not well understood.(161)

While the effects of tensional loading on a variety of protein expression products in tendon are known(159-168), the effect of mechanical strain on hyaluronic acid production and its receptor, CD44, is uncharacterized in tendons. Do these two important mediators of adhesion formation and tendon healing share a common pathway?

There is some evidence that suggest they might. In fibrocartilage tissue, mechanical strain *has* been shown to increase hyaluronic acid production and several of its upstream mediators.(162) If tendon fibroblasts react in a similar way, it would support the view that the beneficial effects of mechanical strain, the cornerstone of current clinical management of tendon healing and adhesion prevention, are acting at least in part through hyaluronic acid, arguably the most promising biochemical player in the prevention of adhesions. With a better understanding of the biochemistry of hyaluronic acid and mechanical strain, a more elegant solution to the problem of adhesions may one day be in hand.

### **Purpose**

The objective of this study was to isolate rat tendon fibroblasts (tenocytes) and to investigate the effect of cyclical mechanical strain on matrix component production, specifically hyaluronic acid (HA), its receptor (CD44) and total glycosaminoglycan (GAG) production.

## **Materials and Methods**

### *Note on individual performance of research tasks:*

All actual labor for the project including, but not limited to, all procedures, cell culture work, animal cell and tissue harvesting, all methods and experiments, generation of data, analysis of data, production of reagents, design and selection of assay systems, use of assay systems including ELISA, western blots, and colorimetric dye-binding assays, preparation of cell lysates, and antibody utilization were performed personally by Richard Crockett. In addition, Richard Crockett was personally responsible for the use, configuration, and repair of the FlexCell unit, as well as the selection, evaluation, procurement and price negotiation of all required reagents and materials, budget management, and project timeline planning.

### Certain elements of the research relied on personnel other than Richard Crockett:

Training of Richard Crockett in the use of western blot technique and photographic film use was performed by Dr. Michael Centrella.

Training of Richard Crockett in surgical dissection techniques for rat tail tendons was performed by Dr. J. Grant Thomson.

Training of Richard Crockett in the culture of tenocytes was performed by Dr. Martijn van Griensven in Hannover, Germany.

Certain elements of the conceptual and pragmatic research plan relied heavily on the creative input of personnel other than Richard Crockett:

Development of laboratory strategies to efficiently obtain data involved discussions with Dr. Thomas McCarthy and Dr. Michael Centrella.

Development of strategies involved in design of the strain protocol and overall research goals involved discussions with Dr. J. Grant Thomson.

Methods Summary:

The research involved several stages, which are explained in detail below. Briefly, rat tail tendons were surgically dissected and the tendon fibroblasts were removed and cultured. After two passages, these cells were transferred to flexible membrane culture dishes and assigned to an experimental or control group. The experimental group was exposed to a cyclical mechanical strain, while the control (or static) group was not. The cells were then analyzed for changes in three targets of interest: hyaluronic acid, the receptor for hyaluronic acid (CD44), and total glycosaminoglycan content.

*Harvest of Tendon Cells:*

The tendon fascicles were harvested from 12 adult Sprague-Dawley rats (Charles River Laboratory; Wilmington, MA). All animal care complied with the IACUC and NIH protocols for the care and use of laboratory animals. The rats were sacrificed with an overdose of isoflurane and cervical dislocation. The tail and hindquarters were

immersed in 95% ethanol then thoroughly rinsed in sterile saline. After transecting the tail, the skin was removed and discarded. The transected tail was transferred to a 150mm Petri dish (Corning; Corning, NY) and using sterile technique, tendon fascicles from dorsal, ventral, and lateral tendon groups in the anterior two-thirds of the tail were removed. After the paratenon was removed, the tissue was divided into 3-5mm<sup>2</sup> samples and placed in 60mm Petri dishes (Corning). For each piece of tissue, a small area of the dish was abraded with a number 11 scalpel blade. The tissue was allowed to partially air dry for several minutes in a sterile laminar flow hood in order to increase adherence. 3-5mL of culture medium was then slowly added. The culture medium consisted of Dulbecco's modified Eagle's Medium (DMEM) with Ham's F-12 (1:1) and L-glutamine (Gibco; Grand Island, NY) supplemented with 10% fetal calf serum (Gibco). Penicillin/streptomycin species (100 U/mL) was also added to the medium.

#### *Culture of Tendon Cells:*

In a 37°C non-moving 5% CO<sub>2</sub> incubator, the tissue was left undisturbed for 2-3 days. Tenocytes migrated from the clumps of tendon tissue. After tenocyte growth reached 75-80% confluence, the tendon tissue was removed. The medium was removed by suction and the dish was bathed in 1-2mL 0.25% trypsin with EDTA (Gibco) at 37°C for 10 minutes. The trypsonized cells were placed in a 15 mL Falcon tube to which culture medium was added. The cells were then collected by sedimentation, resuspended in culture medium and plated in the culture medium on 25cm<sup>2</sup> filter-top culture flasks (Nunc; Roskilde, Denmark). The cells were observed daily under an inverted phase-contrast microscope (Olympus; Melville, NY). Medium was changed every other day.

When the cells had reached 75-80% confluence, the medium was discarded and the cells were freed with trypsin as described above. Cells were then split and plated onto three 80cm<sup>2</sup> filter-top culture flasks. When these cells had reached 75-80% confluence, they were plated on Bioflex® plates (Flexcell International; McKeesport, PA); culture dishes containing a flexible silastic membrane bottom. These culture dishes contained 6 wells, with each well bottom consisting of a flexible membrane which had been previously coated with fibronectin to facilitate fibroblast cell adhesion.

Cultured cells were plated on the same type of Bioflex plate, irrespective of group (strained or static). Only after the cells had been cultured and were ready for the strain protocol were the culture dishes assigned in a random fashion to either the strained or static group. This was done to prevent any selection bias based on cell line. Cell density was determined using a hemocytometer and cells were plated at approximately 10,000 cells/57.75cm<sup>2</sup> Bioflex plate.

*Strain Protocol:*

The Flexercell® Strain Unit applies cyclic strain to cells cultured to confluence on flexible membranes (Flexcell International). The unit exerts a vacuum pressure on a silastic membrane at the bottom of each 6-well plate that has been pre-coated with fibronectin. The unit fits inside a standard CO<sub>2</sub> incubator and the duration, frequency, and magnitude of stress can be adjusted. A 5% change in the size of the membrane approximates a 2% change in the length of cells on the membranes; similar to the amount of elongation collagen experiences under physiologic conditions.(161) To select an appropriate strain regimen, we attempted to mimic active rehabilitation protocols, which

involve periods of repeated strain followed by comparatively longer periods of rest and recovery. Previous studies have shown that the magnitude and duration of strain(163), as well as the length of the rest between strain periods are important parameters(164), but the frequency of strain is less important.(163) Accordingly, cells were strained at 1 Hz for five minutes and allowed to rest for 55 minutes. This was repeated over the course of 8 hours, after which the cells were allowed to rest for 24 hours before collection so that any translational or post-translational events would have sufficient time to occur. Both the medium and cells were collected for analysis.

Cultured tenocytes were divided into two groups – an experimental group, which was exposed to the mechanical strain regimen, and a control or static group which was not exposed to any strain. Both groups were extracted and cultured in identical conditions at the same time and each replicate was derived from an independent cell lineage. During the strain regimen, strain cells which had been cultured on the Bioflex 6-well plates were placed in the strain unit. Static cells were placed in the same incubator at the same time as the strained cells but were not exposed to any strain.

#### *Hyaluronic acid content*

Hyaluronic acid concentration was measured using an enzyme linked protein assay (Corgenix; Westminster, CO). Using a capture molecule called Hyaluronic Acid Binding Protein (HABP) which is bound to horseradish peroxidase (HRP), the assay is visualized with a chromogenic substrate and measured in optical density with a spectrophotometer. Because hyaluronic acid is primarily secreted into the extracellular

tissue matrix, we compared the concentration in the media from strained cells to that found in the media of static cells.

#### *CD44 receptor expression*

Immunolabelling for CD44 was performed using a polyclonal rabbit antibody to CD44 (Santa Cruz Biotechnology; Santa Cruz, CA). After the medium had been removed from the Bioflex plates, the cellular layer was collected and homogenized using a sonicator. The homogenate was normalized for total protein content and separated by charge to mass ratio using SDS polyacrylamide gel electrophoresis. The proteins were electroblotted onto PolyScreen polyvinylidene difluoride transfer membrane (PerkinElmer Life Sciences, Wellesley, MA). The western blot was then probed using primary antibody (as described above) in TBST with milk at 20°C overnight. The secondary antibody was goat, anti-rabbit conjugated to HRP (Santa Cruz) and immunocomplex formation was visualized with chemiluminescence and exposure to photographic film.

#### *Total glycosaminoglycan (GAG) content*

To measure the amount of glycosaminoglycan (GAG) present in the medium, a colorimetric assay was used which involves the binding of a dye (dimethyl-methylene blue) to GAGs (Biocolor Assays; Newtownabbey, Northern Ireland). A dye solution which includes DMB dye and an inorganic solvent are added to the sample. The dye is in excess and binds with GAGs, precipitating out of solution. After centrifugation, the resulting pellet contains the dye-GAG complex, and the supernatant contains unbound

DMB dye and solvent. Resuspension of the pellet in propanol to disassociate the dye from the GAGs allows the concentration of solubilized dye to be read on a spectrophotometer.

### **Data Analysis**

An unpaired two-tailed Student's t-test was used to compare strained and static groups. A value of  $p < 0.05$  was considered significant. Western blots were optically scanned and analyzed using ImageJ (NIH; Bethesda, MD).

## **Results**

During the harvest of the tendon fibroblasts, 12 animals were dissected and the tendons were removed. Two additional animals were used solely for practice of the tissue harvest and culture and were not part of the experiment. Although it was initially challenging to separate the tendon from the surrounding tissue (including the paratenon), once the technical elements were refined the process became straightforward. Each animal yielded a sufficient quantity of tendon tissue such that each control and experimental cell line consisted of tenocytes with a distinct cell lineage from the others. These were cultured in parallel, under identical conditions. One culture developed a fungal infection before assignment to either the experimental or control group and was discarded.

### *Hyaluronic Acid Content*

Since hyaluronic acid is produced by fibroblasts and found primarily in the extracellular environment, the medium in which the cells were cultured was analyzed quantitatively for hyaluronic acid content. The removed media were normalized for total protein content before analysis. Using an ELISA assay, 44 medium samples (22 experimental, 22 control) were analyzed. The medium of tenocytes which had experienced cyclical mechanical strain had a greater concentration of hyaluronic acid ( $1528 \pm 58$  ng/mL (mean  $\pm$  SEM)) compared with the medium of tenocytes that had not been subject to strain ( $730 \pm 27$  ng/mL). This constitutes nearly a two fold increase in hyaluronic acid content ( $p < .0001$ ). (See Table 1).


**Table 1**

Group	Assay		
	Hyaluronic Acid <i>ng/mL</i>	CD44 Receptor <i>O.D.</i>	Glycosaminoglycans <i>μg/mL</i>
Strained	1528 +/- 58	25525 +/- 2461	10.2 +/- 0.6
Static	793 +/- 27	15454 +/- 1443	15.3 +/- 3.0
p value	< .0001	0.024	0.103

Comparisons between strained and static cells for hyaluronic acid, CD44, and glycosaminoglycan content. Values are mean  $\pm$  standard error of the mean.

#### *CD44 receptor expression*

Hyaluronic acid as a signaling molecule acts primarily through its receptor, CD44. To assess whether there were changes in the population of the CD44 receptor, the cell membranes of the tenocytes were analyzed for the presence of CD44. To allow for the transcription and membrane translocation of the receptors, a delay of 24 hours between the end of the mechanical strain regimen and collection was instituted. Using a western blot analysis, 6 (3 experimental, 3 control) cellular homogenate samples from independent cell lineages were analyzed for expression of CD44. (See figure 1). Tenocytes under mechanical strain increased their concentration of CD44 by 62.5%, with tenocytes exposed to strain having an optical density of  $26 \cdot 10^3 \pm 2 \cdot 10^3$  compared with  $15 \cdot 10^3 \pm 1 \cdot 10^3$  in controls ( $p < .05$ ). (See table 1).

**Figure 1**

Inset of western blot for CD44. Lanes 1 through 3 represent the increased signal in experimental cells exposed to cyclic strain compared to control lanes 4 through 6 which shows CD44 production in static cells.

*Total glycosaminoglycan content*

The total amount of glycosaminoglycan produced by tenocytes in both the strained and static groups was measured to examine what effect mechanical strain had on the production of these structural polysaccharides. Using a commercially available colorimetric dye-binding assay, 44 medium samples (22 experimental, 22 control) were analyzed for content of glycosaminoglycans. No significant differences were detected in glycosaminoglycan content between the media of tenocytes which had been cyclically strained versus cells which were not, with strained cells producing  $10.2 \pm 0.6 \mu\text{g/mL}$  and controls producing  $15.3 \pm 3.0 \mu\text{g/mL}$  ( $p = 0.103$ ). (See Table 1).

## **Discussion**

Proper tendon healing after repair requires a balance between appropriate fibrous connective tissue formation to provide tensile strength and the formation of excessive scar tissue, fibrosis, which impedes the smooth gliding of structures and leads to adhesions and impairment of function. Mechanical strain has been shown important in this collagen production and the linear organization of collagen fibrils(54). Its role both in animal models and clinical settings is widely accepted. The active rehabilitation protocols used for hand surgery patients currently employed are designed to tread between these two extremes: too little activity or strain can result in the formation of adhesions, while excessive strain can result in rupture at the site of tenorrhaphy.

Understanding the effects of mechanical strain on tendon fibroblasts is important in finding adjunct medical therapies which would mitigate fibrosis and adhesion formation without weakening tendon repair. One such candidate is hyaluronic acid—a high molecular weight polysaccharide found throughout the body that provides lubrication in joints and the smooth gliding of tendons. More recently, research has shown that in addition to its structural and lubricating role, hyaluronic acid as an important signaling molecule for cell motility, adhesion, and proliferation.

Hyaluronic acid has been shown to be beneficial in preventing or decreasing adhesions in animal models including chicken(142, 144, 149), rabbit(137, 143, 148, 152, 158), canine(136), horse(141), and primate.(150) Unfortunately, this success in animal models has not translated into clinical applications.(42, 153, 154) This disappointing gap suggests that a closer look at hyaluronic acid may be warranted.

Our study examined whether mechanical stress might directly influence hyaluronic acid regulation. The model we chose was one in which rat tail tenocytes were exposed to cyclic strain *in vitro*. One question that this raises is whether this is a valid model for examining the effects on tendon fibroblasts.

Dissociated cells and tendon explants do appear to be good models for examining the effects of mechanical strain on tenocytes. Dissociated tenocytes bond well to fibronectin as demonstrated in biomaterial studies which examine adhesion strength.(165) Tendon explants and dissociated fibroblasts grown in culture retain their ability to respond to mechanical stress.(111, 161, 166-171) Moreover, the effects seen correlate well with the effects of mechanical strain seen *in vivo*.(172-176)

Part of the difficulty in teasing apart the role of hyaluronic acid is that it seems to behave differently based on its molecular weight, concentration, and the presence of other growth factors. In fact, despite evidence that hyaluronic acid is a key molecule in tissue healing, cell migration, and adhesion formation, we know little about how it is regulated.

Probably the most important regulator of hyaluronic acid is hyaluronic acid synthase (HAS), particularly the second isoform, Has2(177, 178). In a study that examined wound healing with cell lines that expressed sense and anti-sense Has2, the increased cell migration and faster healing seen with sense Has2 was absent in cells expressing anti-sense Has2. The anti-sense Has2 cells were also associated with an increase in adhesive plaques. Exogenous hyaluronate was not able to overcome the deficiency, suggesting that activation of Has2, more than the quantity of hyaluronic acid, is the key determiner of how adhesions are formed and regulated by the hyaluronan pathway.(179)

This raises the question: if hyaluronic acid is beneficial to tendon healing, would inhibiting the enzymes which degrade it (hyaluronidases) be beneficial as well? The answer in tendons has not been explored. In intra-peritoneal adhesions however, where some data suggest that hyaluronic acid is helpful in reducing the adhesion formation(180), the effect has been examined(181, 182) and some researchers have seen a benefit.(183)

In sum, it seems that hyaluronic acid has more than one role as a signaling molecule. At a molecular level, hyaluronidase can suppress the rapid interactions that occur early in cell adhesion formation. When a small amount of hyaluronic acid exists, it can serve to facilitate cell-matrix adhesion, whereas excess hyaluronic acid strongly inhibits cell adhesion.(184) Thus, it seems that hyaluronic acid may help to allow cells to interact, move and bind together, while at the same time, larger amounts may help prevent adhesion formation.

This study showed a significant increase in both hyaluronic acid and its receptor (CD44) when tenocytes were exposed to cyclical mechanical strain *in vitro*. Our study is the first to examine the effects of mechanical strain on the regulation of hyaluronic acid in tendons. It is clear from our study that mechanical strain, which is an integral part of the repair process, plays a role in the regulation of tendon fibroblast hyaluronic acid production. How that translates into adhesion formation still needs to be explored, however.

When we examined the total glycosaminoglycan content, it was not found to be significantly altered by mechanical strain, though other investigators have shown previously that many stimuli including *in vitro* culture alone can alter total and sulphated

GAG content(111). The increase in hyaluronic acid, but not total glycosaminoglycan content, may reflect the preferential effect of mechanical strain on the hyaluronan pathway. Another possibility is that we did not observe any change in glycosaminoglycan content in response to mechanical strain because of limitations of the assay, which performed poorly even during calibration of standards. Additionally, the assay we performed examined the presence of glycosaminoglycans in the supernatant. The predominance of glycosaminoglycans may have been intimately associated with the cell membranes and not readily present in the supernatant.

Hyaluronic acid is a promising target for mitigating adhesion formation. Unfortunately, the clinical applications to date related to post-operative tendon repair have been disappointing(153, 154). In addition to the complexities of its regulation and mechanisms of action, use of hyaluronic acid has been challenged by its rapid catabolism and dilution in the body.

Newer, derivatized formulations of HA show longer persistence and less friction in tendons after repair(185) and have been used for a variety of purposes(186) from coating metallic artificial joints(187) to forming biologically compatible hydrogels.(188) Other researchers have created cross-linked hyaluronic acid films which contain DNA encoding hyaluronan synthase 2 (HAS2). The goal of these biofilms is to provide a physical barrier while at the same time delivering hyaluronan and upregulating HAS. The films are subject to the same catabolism by hyaluronidases as native hyaluronic acid, and as such subject to breakdown, but cross-linked hyaluronic acid-collagen sheets which are resistant to hyaluronidase over several weeks have been developed.(189) Hopefully, these promising discoveries will translate into viable clinical applications.

The future role of hyaluronic acid as an adjunct medical therapy to prevent the formation of adhesions after tendon surgery is unclear. The current challenges in understanding how hyaluronic acid fits into the larger process of wound healing are interesting avenues of research, however, and as the details of its regulation and mechanisms of action are revealed, perhaps a clearer path to preventing tendon adhesions will be as well.

## References

1. Manske, P.R., History of flexor tendon repair. *Hand Clin*, 2005. 21(2): 123-7.
2. Strickland, J.W., Flexor tendon injuries. Part 1. Anatomy, physiology, biomechanics, healing, and adhesion formation around a repaired tendon. *Orthop Rev*, 1986. 15(10): 632-45.
3. Potenza, A.D., Tendon healing within the flexor digital sheath in the dog. *J Bone Joint Surg Am*, 1962. 44-A: 49-64.
4. Gelberman, R.H. and P.R. Manske, Factors influencing flexor tendon adhesions. *Hand Clin*, 1985. 1(1): 35-42.
5. Hagberg, L., O. Wik, and B. Gerdin, Determination of biomechanical characteristics of restrictive adhesions and of functional impairment after flexor tendon surgery: a methodological study of rabbits. *J Biomech*, 1991. 24(10): 935-42.
6. Masuda, K., et al., Biochemical analysis of collagen in adhesive tissues formed after digital flexor tendon injuries. *J Orthop Sci*, 2002. 7(6): 665-71.
7. Ragoowansi, R., et al., Differences in morphology, cytoskeletal architecture and protease production between zone II tendon and synovial fibroblasts in vitro. *J Hand Surg [Br]*, 2003. 28(5): 465-70.
8. Belmahi, A., N.E. Gharib, and S. El Mazouz, [The protected tendinous graft in zone 2 or fibroblast trap]. *Chir Main*, 2004. 23(3): 142-8.
9. Frykman, E., S. Jacobsson, and B. Widenfalk, Fibrin sealant in prevention of flexor tendon adhesions: an experimental study in the rabbit. *J Hand Surg [Am]*, 1993. 18(1): 68-75.
10. Tan, V., et al., Interosseous-lumbrical adhesions of the hand: contribution of magnetic resonance imaging to diagnosis and treatment planning. *J Hand Surg [Am]*, 2002. 27(4): 639-43.
11. Langley, P.L., Iontophoresis to aid in releasing tendon adhesions. Suggestion from the field. *Phys Ther*, 1984. 64(9): 1395.
12. Robotti, E., et al., The effect of pulsed electromagnetic fields on flexor tendon healing in chickens. *J Hand Surg [Br]*, 1999. 24(1): 56-8.
13. Greenough, C.G., The effect of pulsed electromagnetic fields on flexor tendon healing in the rabbit. *J Hand Surg [Br]*, 1996. 21(6): 808-12.
14. Healy, C., et al., Postoperative stiffness and adhesion formation around repaired and immobilized Achilles tenotomies are prevented using a model of heat shock protein induction. *J Surg Res*, 2004. 120(2): 225-9.
15. Mulhall, K.J., et al., Thermal preconditioning prevents peritendinous adhesions and inflammation. *Clin Orthop Relat Res*, 2002(405): 258-66.
16. Szabo, R.M. and E. Younger, Effects of indomethacin on adhesion formation after repair of zone II tendon lacerations in the rabbit. *J Hand Surg [Am]*, 1990. 15(3): 480-3.

17. Carlstedt, C.A., Mechanical and chemical factors in tendon healing. Effects of indomethacin and surgery in the rabbit. *Acta Orthop Scand Suppl*, 1987. 224: 1-75.
18. Carlstedt, C.A., K. Madsen, and T. Wredmark, The influence of indomethacin on biomechanical and biochemical properties of the plantaris longus tendon in the rabbit. *Arch Orthop Trauma Surg*, 1987. 106(3): 157-60.
19. Carlstedt, C.A., K. Madsen, and T. Wredmark, The influence of indomethacin on tendon healing. A biomechanical and biochemical study. *Arch Orthop Trauma Surg*, 1986. 105(6): 332-6.
20. Kulick, M.I., S. Smith, and K. Hadler, Oral ibuprofen: evaluation of its effect on peritendinous adhesions and the breaking strength of a tenorrhaphy. *J Hand Surg [Am]*, 1986. 11(1): 110-20.
21. Kulick, M.I., et al., Injectable ibuprofen: preliminary evaluation of its ability to decrease peritendinous adhesions. *Ann Plast Surg*, 1984. 13(6): 459-67.
22. Sullo, A., et al., The effects of prolonged peritendinous administration of PGE1 to the rat Achilles tendon: a possible animal model of chronic Achilles tendinopathy. *J Orthop Sci*, 2001. 6(4): 349-57.
23. Ketchum, L.D., Effects of triamcinolone on tendon healing and function. A laboratory study. *Plast Reconstr Surg*, 1971. 47(5): 471-82.
24. Kapetanos, G., The effect of the local corticosteroids on the healing and biomechanical properties of the partially injured tendon. *Clin Orthop Relat Res*, 1982(163): 170-9.
25. Douglas, L.G., S.H. Jackson, and W.K. Lindsay, The effects of dexamethasone, norethandrolone, promethazine and a tension-relieving procedure on collagen synthesis in healing flexor tendons as estimated by tritiated proline uptake studies. *Can J Surg*, 1967. 10(1): 36-46.
26. Walker, F.G., S.H. Bensley, and W.K. Lindsay, The effects of an antihistamine (promethazine) on the reaction of tendons to trauma (a histological study). *Acta Chir Plast*, 1965. 7(3): 171-86.
27. Lane, J.M., J. Black, and F.W. Bora, Jr., Gliding function following flexor-tendon injury. A biomechanical study of rat tendon function. *J Bone Joint Surg Am*, 1976. 58(7): 985-90.
28. Akali, A., et al., Decrease in adhesion formation by a single application of 5-fluorouracil after flexor tendon injury. *Plast Reconstr Surg*, 1999. 103(1): 151-8.
29. Khan, U., et al., Modulation of the formation of adhesions during the healing of injured tendons. *J Bone Joint Surg Br*, 2000. 82(7): 1054-8.
30. Moran, S.L., et al., Effects of 5-fluorouracil on flexor tendon repair. *J Hand Surg [Am]*, 2000. 25(2): 242-51.
31. Cerovac, S., et al., Early breaking strength of repaired flexor tendon treated with 5-fluorouracil. *J Hand Surg [Br]*, 2001. 26(3): 220-3.
32. Ragoowansi, R., et al., Reduction in matrix metalloproteinase production by tendon and synovial fibroblasts after a single exposure to 5-fluorouracil. *Br J Plast Surg*, 2001. 54(4): 283-7.
33. Komurcu, M., et al., Reduction of restrictive adhesions by local aprotinin application and primary sheath repair in surgically traumatized flexor tendons of the rabbit. *J Hand Surg [Am]*, 1997. 22(5): 826-32.

34. Nyska, M., et al., Topically applied halofuginone, an inhibitor of collagen type I transcription, reduces peritendinous fibrous adhesions following surgery. *Connect Tissue Res*, 1996. 34(2): 97-103.
35. Speer, D.P., S. Feldman, and M. Chvapil, The control of peritendinous adhesions using topical beta-aminopropionitrile base. *J Surg Res*, 1985. 38(3): 252-7.
36. Porat, S., et al., Increased collagenolytic activity in severed and sutured tendons following topical application of exogenous collagen in chickens. *J Orthop Res*, 1985. 3(1): 43-8.
37. Porat, S., M. Rousso, and S. Shoshan, Improvement of gliding function of flexor tendons by topically applied enriched collagen solution. *J Bone Joint Surg Br*, 1980. 62-B(2): 208-13.
38. Nyska, M., et al., Decreased adhesion formation in flexor tendons by topical application of enriched collagen solution--a histological study. *Arch Orthop Trauma Surg*, 1987. 106(3): 192-4.
39. Hanff, G. and L. Hagberg, Prevention of restrictive adhesions with expanded polytetrafluoroethylene diffusible membrane following flexor tendon repair: an experimental study in rabbits. *J Hand Surg [Am]*, 1998. 23(4): 658-64.
40. Kobayashi, M., J. Toguchida, and M. Oka, Development of polyvinyl alcohol-hydrogel (PVA-H) shields with a high water content for tendon injury repair. *J Hand Surg [Br]*, 2001. 26(5): 436-40.
41. Gudemez, E., et al., Chondroitin sulfate-coated polyhydroxyethyl methacrylate membrane prevents adhesion in full-thickness tendon tears of rabbits. *J Hand Surg [Am]*, 2002. 27(2): 293-306.
42. Strickland, J.W., Development of flexor tendon surgery: twenty-five years of progress. *J Hand Surg [Am]*, 2000. 25(2): 214-35.
43. Siddiqi, N.A., et al., Effects of hydroxyapatite and alumina sheaths on postoperative peritendinous adhesions in chickens. *J Appl Biomater*, 1995. 6(1): 43-53.
44. Smith, D.J., Jr., et al., Use of glutaraldehyde stabilized mammalian pericardium in hand surgery. *Ann Chir Main*, 1988. 7(1): 54-7.
45. Stark, H.H., et al., The use of paratenon, polyethylene film, or silastic sheeting to prevent restricting adhesions to tendons in the hand. *J Bone Joint Surg Am*, 1977. 59(7): 908-13.
46. Golash, A., et al., Efficacy of ADCON-T/N after primary flexor tendon repair in Zone II: a controlled clinical trial. *J Hand Surg [Br]*, 2003. 28(2): 113-5.
47. Mentzel, M., et al., The effectiveness of ADCON-T/N, a new anti-adhesion barrier gel, in fresh divisions of the flexor tendons in Zone II. *J Hand Surg [Br]*, 2000. 25(6): 590-2.
48. Silfverskiold, K.L. and E.J. May, Flexor tendon repair in zone II with a new suture technique and an early mobilization program combining passive and active flexion. *J Hand Surg [Am]*, 1994. 19(1): 53-60.
49. Peck, F.H., et al., A comparative study of two methods of controlled mobilization of flexor tendon repairs in zone 2. *J Hand Surg [Br]*, 1998. 23(1): 41-5.
50. Harris, S.B., et al., The aetiology of acute rupture of flexor tendon repairs in zones 1 and 2 of the fingers during early mobilization. *J Hand Surg [Br]*, 1999. 24(3): 275-80.

51. Lister, G.D., et al., Primary flexor tendon repair followed by immediate controlled mobilization. *J Hand Surg [Am]*, 1977. 2(6): 441-51.
52. Becker, H., F. Orak, and E. Duponselle, Early active motion following a beveled technique of flexor tendon repair: report on fifty cases. *J Hand Surg [Am]*, 1979. 4(5): 454-60.
53. Strickland, J.W., Results of flexor tendon surgery in zone II. *Hand Clin*, 1985. 1(1): 167-79.
54. Strickland, J.W. and S.V. Glogovac, Digital function following flexor tendon repair in Zone II: A comparison of immobilization and controlled passive motion techniques. *J Hand Surg [Am]*, 1980. 5(6): 537-43.
55. Gault, D.T., Reduction of grip strength, finger flexion pressure, finger pinch pressure and key pinch following flexor tendon repair. *J Hand Surg [Br]*, 1987. 12(2): 182-4.
56. Singer, M. and S. Maloon, Flexor tendon injuries: the results of primary repair. *J Hand Surg [Br]*, 1988. 13(3): 269-72.
57. Chow, J.A., et al., Controlled motion rehabilitation after flexor tendon repair and grafting. A multi-centre study. *J Bone Joint Surg Br*, 1988. 70(4): 591-5.
58. Small, J.O., M.D. Brennen, and J. Colville, Early active mobilisation following flexor tendon repair in zone 2. *J Hand Surg [Br]*, 1989. 14(4): 383-91.
59. Cullen, K.W., et al., Flexor tendon repair in zone 2 followed by controlled active mobilisation. *J Hand Surg [Br]*, 1989. 14(4): 392-5.
60. Savage, R. and G. Risitano, Flexor tendon repair using a "six strand" method of repair and early active mobilisation. *J Hand Surg [Br]*, 1989. 14(4): 396-9.
61. May, E.J., K.L. Silfverskiold, and C.J. Sollerman, Controlled mobilization after flexor tendon repair in zone II: a prospective comparison of three methods. *J Hand Surg [Am]*, 1992. 17(5): 942-52.
62. Bainbridge, L.C., et al., A comparison of post-operative mobilization of flexor tendon repairs with "passive flexion-active extension" and "controlled active motion" techniques. *J Hand Surg [Br]*, 1994. 19(4): 517-21.
63. Elliot, D., et al., The rupture rate of acute flexor tendon repairs mobilized by the controlled active motion regimen. *J Hand Surg [Br]*, 1994. 19(5): 607-12.
64. Baktir, A., et al., Flexor tendon repair in zone 2 followed by early active mobilization. *J Hand Surg [Br]*, 1996. 21(5): 624-8.
65. Kitsis, C.K., et al., Controlled active motion following primary flexor tendon repair: a prospective study over 9 years. *J Hand Surg [Br]*, 1998. 23(3): 344-9.
66. Aoki, M., et al., Biomechanical and histologic characteristics of canine flexor tendon repair using early postoperative mobilization. *J Hand Surg [Am]*, 1997. 22(1): 107-14.
67. Pruitt, D.L., et al., Cyclic stress testing after in vivo healing of canine flexor tendon lacerations. *J Hand Surg [Am]*, 1996. 21(6): 974-7.
68. Wada, A., et al., Comparison of postoperative early active mobilization and immobilization in vivo utilising a four-strand flexor tendon repair. *J Hand Surg [Br]*, 2001. 26(4): 301-6.
69. Feehan, L.M. and J.G. Beauchene, Early tensile properties of healing chicken flexor tendons: early controlled passive motion versus postoperative immobilization. *J Hand Surg [Am]*, 1990. 15(1): 63-8.

70. Gelberman, R.H., et al., The excursion and deformation of repaired flexor tendons treated with protected early motion. *J Hand Surg [Am]*, 1986. 11(1): 106-10.
71. Gelberman, R.H., et al., Effects of early intermittent passive mobilization on healing canine flexor tendons. *J Hand Surg [Am]*, 1982. 7(2): 170-5.
72. Gelberman, R.H., et al., Flexor tendon repair. *J Orthop Res*, 1986. 4(1): 119-28.
73. Takai, S., et al., The effects of frequency and duration of controlled passive mobilization on tendon healing. *J Orthop Res*, 1991. 9(5): 705-13.
74. Gelberman, R.H., et al., Influences of the protected passive mobilization interval on flexor tendon healing. A prospective randomized clinical study. *Clin Orthop Relat Res*, 1991(264): 189-96.
75. Evans, R.B., Clinical application of controlled stress to the healing extensor tendon: a review of 112 cases. *Phys Ther*, 1989. 69(12): 1041-9.
76. Evans, R.B. and D.E. Thompson, The application of force to the healing tendon. *J Hand Ther*, 1993. 6(4): 266-84.
77. Chao, R.P., et al., Early passive mobilization after digital nerve repair and grafting in a fresh cadaver. *Plast Reconstr Surg*, 2001. 108(2): 386-91.
78. Woo, S.L., et al., The importance of controlled passive mobilization on flexor tendon healing. A biomechanical study. *Acta Orthop Scand*, 1981. 52(6): 615-22.
79. Bunker, T.D., B. Potter, and N.J. Barton, Continuous passive motion following flexor tendon repair. *J Hand Surg [Br]*, 1989. 14(4): 406-11.
80. Tran, H.N., et al., In vitro cyclic tensile testing of combined peripheral and core flexor tenorrhaphy suture techniques. *J Hand Surg [Am]*, 2002. 27(3): 518-24.
81. Zhao, C., et al., Remodeling of the gliding surface after flexor tendon repair in a canine model in vivo. *J Orthop Res*, 2002. 20(4): 857-62.
82. Momose, T., et al., Surface modification of extrasynovial tendon by chemically modified hyaluronic acid coating. *J Biomed Mater Res*, 2002. 59(2): 219-24.
83. Matthews, P. and H. Richards, Factors in the adherence of flexor tendon after repair: an experimental study in the rabbit. *J Bone Joint Surg Br*, 1976. 58(2): 230-6.
84. Banes, A.J., et al., Mechanical load stimulates expression of novel genes in vivo and in vitro in avian flexor tendon cells. *Osteoarthritis Cartilage*, 1999. 7(1): 141-53.
85. Orhan, Z., et al., The effect of extracorporeal shock waves on a rat model of injury to tendo Achillis. A histological and biomechanical study. *J Bone Joint Surg Br*, 2004. 86(4): 613-8.
86. Tanaka, H., et al., Effect of cyclic tension on lacerated flexor tendons in vitro. *J Hand Surg [Am]*, 1995. 20(3): 467-73.
87. Mass, D.P., et al., Effects of constant mechanical tension on the healing of rabbit flexor tendons. *Clin Orthop Relat Res*, 1993(296): 301-6.
88. Wray, R.C., Jr., et al., Effect of continuous load on the mechanical properties of tendon adhesions. *Hand*, 1981. 13(1): 92-6.
89. Kubota, H., et al., Effect of motion and tension on injured flexor tendons in chickens. *J Hand Surg [Am]*, 1996. 21(3): 456-63.
90. Silva, M.J., et al., Effects of increased in vivo excursion on digital range of motion and tendon strength following flexor tendon repair. *J Orthop Res*, 1999. 17(5): 777-83.

91. Fong, K.D., et al., Microarray analysis of mechanical shear effects on flexor tendon cells. *Plast Reconstr Surg*, 2005. 116(5): 1393-404; discussion 1405-6.
92. Abrahamsson, S.O., G. Lundborg, and L.S. Lohmander, Tendon healing in vivo. An experimental model. *Scand J Plast Reconstr Surg Hand Surg*, 1989. 23(3): 199-205.
93. Abrahamsson, S.O., G. Lundborg, and L.S. Lohmander, Recombinant human insulin-like growth factor-I stimulates in vitro matrix synthesis and cell proliferation in rabbit flexor tendon. *J Orthop Res*, 1991. 9(4): 495-502.
94. Abrahamsson, S.O., G. Lundborg, and L.S. Lohmander, Long-term explant culture of rabbit flexor tendon: effects of recombinant human insulin-like growth factor-I and serum on matrix metabolism. *J Orthop Res*, 1991. 9(4): 503-15.
95. Zhang, A.Y., et al., Inhibition of TGF-beta-induced collagen production in rabbit flexor tendons. *J Hand Surg [Am]*, 2004. 29(2): 230-5.
96. Klein, M.B., et al., Flexor tendon healing in vitro: effects of TGF-beta on tendon cell collagen production. *J Hand Surg [Am]*, 2002. 27(4): 615-20.
97. Chang, J., et al., Studies in flexor tendon wound healing: neutralizing antibody to TGF-beta1 increases postoperative range of motion. *Plast Reconstr Surg*, 2000. 105(1): 148-55.
98. Chang, J., et al., Gene expression of transforming growth factor beta-1 in rabbit zone II flexor tendon wound healing: evidence for dual mechanisms of repair. *Plast Reconstr Surg*, 1997. 100(4): 937-44.
99. Tsubone, T., et al., Expression of growth factors in canine flexor tendon after laceration in vivo. *Ann Plast Surg*, 2004. 53(4): 393-7.
100. Ngo, M., et al., Differential expression of transforming growth factor-beta receptors in a rabbit zone II flexor tendon wound healing model. *Plast Reconstr Surg*, 2001. 108(5): 1260-7.
101. Gelberman, R.H., et al., Angiogenesis in healing autogenous flexor-tendon grafts. *J Bone Joint Surg Am*, 1992. 74(8): 1207-16.
102. Ditsios, K., et al., Neovascularization of the flexor digitorum profundus tendon after avulsion injury: an in vivo canine study. *J Hand Surg [Am]*, 2003. 28(2): 231-6.
103. Gelberman, R.H., et al., The effects of mobilization on the vascularization of healing flexor tendons in dogs. *Clin Orthop Relat Res*, 1980(153): 283-9.
104. Darmani, H., J.C. Crossan, and A. Curtis, Single dose of inducible nitric oxide synthase inhibitor induces prolonged inflammatory cell accumulation and fibrosis around injured tendon and synovium. *Mediators Inflamm*, 2004. 13(3): 157-64.
105. Klein, M.B., et al., Flexor tendon wound healing in vitro: the effect of lactate on tendon cell proliferation and collagen production. *J Hand Surg [Am]*, 2001. 26(5): 847-54.
106. Mehta, V., et al., Characterization of adenovirus-mediated gene transfer in rabbit flexor tendons. *J Hand Surg [Am]*, 2005. 30(1): 136-41.
107. Shimomura, T., et al., Antisense oligonucleotides reduce synthesis of procollagen alpha1 (V) chain in human patellar tendon fibroblasts: potential application in healing ligaments and tendons. *Connect Tissue Res*, 2003. 44(3-4): 167-72.
108. Dykxj, D. and K.T. Jules, The clinical anatomy of tendons. *J Am Podiatr Med Assoc*, 1991. 81(7): 358-65.

109. Vailas, A.C., et al., Physical activity and hypophysectomy on the aerobic capacity of ligaments and tendons. *J Appl Physiol*, 1978. 44(4): 542-6.
110. Williams, J.G., Achilles tendon lesions in sport. *Sports Med*, 1986. 3(2): 114-35.
111. Slack, C., et al., Changes in the morphology and synthetic activity of cultured rat tail tendon. *Cell Tissue Res*, 1986. 245(2): 359-68.
112. Kastelic, J., A. Galeski, and E. Baer, The multicomposite structure of tendon. *Connect Tissue Res*, 1978. 6(1): 11-23.
113. Wilson, T.W., M.P. Zafuta, and M. Zobitz, A biomechanical analysis of matched bone-patellar tendon-bone and double-looped semitendinosus and gracilis tendon grafts. *Am J Sports Med*, 1999. 27(2): 202-7.
114. Tammi, M.I., A.J. Day, and E.A. Turley, Hyaluronan and homeostasis: a balancing act. *J Biol Chem*, 2002. 277(7): 4581-4.
115. Day, A.J. and G.D. Prestwich, Hyaluronan-binding proteins: tying up the giant. *J Biol Chem*, 2002. 277(7): 4585-8.
116. Olutoye, O.O., et al., Hyaluronic acid inhibits fetal platelet function: implications in scarless healing. *J Pediatr Surg*, 1997. 32(7): 1037-40.
117. Shepard, S., H. Becker, and J.X. Hartmann, Using hyaluronic acid to create a fetal-like environment in vitro. *Ann Plast Surg*, 1996. 36(1): 65-9.
118. Alaish, S.M., et al., Biology of fetal wound healing: hyaluronate receptor expression in fetal fibroblasts. *J Pediatr Surg*, 1994. 29(8): 1040-3.
119. Dillon, P.W., et al., The extracellular matrix of the fetal wound: hyaluronic acid controls lymphocyte adhesion. *J Surg Res*, 1994. 57(1): 170-3.
120. al-Qattan, M.M., et al., Fetal tendon healing: development of an experimental model. *Plast Reconstr Surg*, 1993. 92(6): 1155-60; discussion 1161.
121. Meinert, M., et al., Proteoglycans and hyaluronan in human fetal membranes. *Am J Obstet Gynecol*, 2001. 184(4): 679-85.
122. Luo, J., Z. Yang, and X. Li, [Effect of human acellular amnion membrane on tendon adhesion in rat]. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi*, 2004. 18(5): 431-4.
123. Demirkan, F., et al., The use of amniotic membrane in flexor tendon repair: an experimental model. *Arch Orthop Trauma Surg*, 2002. 122(7): 396-9.
124. Ozgenel, G.Y., The effects of a combination of hyaluronic and amniotic membrane on the formation of peritendinous adhesions after flexor tendon surgery in chickens. *J Bone Joint Surg Br*, 2004. 86(2): 301-7.
125. Ozgenel, G.Y., B. Samli, and M. Ozcan, Effects of human amniotic fluid on peritendinous adhesion formation and tendon healing after flexor tendon surgery in rabbits. *J Hand Surg [Am]*, 2001. 26(2): 332-9.
126. Obara, T., et al., Increased friction of animal joints by experimental degeneration and recovery by addition of hyaluronic acid. *Clin Biomech (Bristol, Avon)*, 1997. 12(4): 246-252.
127. Mabuchi, K., et al., The effect of additive hyaluronic acid on animal joints with experimentally reduced lubricating ability. *J Biomed Mater Res*, 1994. 28(8): 865-70.
128. Noble, P.W., Hyaluronan and its catabolic products in tissue injury and repair. *Matrix Biol*, 2002. 21(1): 25-9.

129. Mabuchi, K., et al., Molecular weight independence of the effect of additive hyaluronic acid on the lubricating characteristics in synovial joints with experimental deterioration. *Clin Biomech (Bristol, Avon)*, 1999. 14(5): 352-6.
130. Collis, L., et al., Rapid hyaluronan uptake is associated with enhanced motility: implications for an intracellular mode of action. *FEBS Lett*, 1998. 440(3): 444-9.
131. Evanko, S.P. and T.N. Wight, Intracellular localization of hyaluronan in proliferating cells. *J Histochem Cytochem*, 1999. 47(10): 1331-42.
132. Aruffo, A., et al., CD44 is the principal cell surface receptor for hyaluronate. *Cell*, 1990. 61(7): 1303-13.
133. Goldstein, L.A., et al., A human lymphocyte homing receptor, the hermes antigen, is related to cartilage proteoglycan core and link proteins. *Cell*, 1989. 56(6): 1063-72.
134. Pure, E. and C.A. Cuff, A crucial role for CD44 in inflammation. *Trends Mol Med*, 2001. 7(5): 213-21.
135. Lesley, J. and R. Hyman, CD44 structure and function. *Front Biosci*, 1998. 3: d616-30.
136. Amiel, D., et al., Hyaluronan in flexor tendon repair. *J Hand Surg [Am]*, 1989. 14(5): 837-43.
137. Thomas, S.C., L.C. Jones, and D.S. Hungerford, Hyaluronic acid and its effect on postoperative adhesions in the rabbit flexor tendon. A preliminary look. *Clin Orthop Relat Res*, 1986(206): 281-9.
138. Akasaka, T., et al., Hyaluronic acid diminishes the resistance to excursion after flexor tendon repair: an in vitro biomechanical study. *J Biomech*, 2005. 38(3): 503-7.
139. Nishida, J., et al., Effect of hyaluronic acid on the excursion resistance of tendon grafts. A biomechanical study in a canine model in vitro. *J Bone Joint Surg Br*, 2004. 86(6): 918-24.
140. Halici, M., et al., Sodium hyaluronate regulating angiogenesis during Achilles tendon healing. *Knee Surg Sports Traumatol Arthrosc*, 2004. 12(6): 562-7.
141. Gaughan, E.M., et al., Effects of sodium hyaluronate on tendon healing and adhesion formation in horses. *Am J Vet Res*, 1991. 52(5): 764-73.
142. Yuzawa, K., [Experimental studies on the healing and restoration of gliding function of the injured digital flexor tendon. Part 9: The use of drugs to prevent adhesion formation of the injured tendon]. *Nippon Seikeigeka Gakkai Zasshi*, 1985. 59(12): 1107-18.
143. Xu, J., Y. Gu, and H. Wang, [Experimental study of sodium hyaluronate products on prevention of tendon adhesion]. *Zhonghua Wai Ke Za Zhi*, 1995. 33(9): 529-31.
144. Chen, Z. and J. Gu, [Experimental research on using macromolecule sodium hyaluronate to prevent flexor tendon adhesion]. *Zhonghua Wai Ke Za Zhi*, 1995. 33(9): 526-8.
145. Tuncay, I., et al., Effects of hyaluronic acid on postoperative adhesion of tendo calcaneus surgery: an experimental study in rats. *J Foot Ankle Surg*, 2002. 41(2): 104-8.
146. Meyers, S.A., et al., Effect of hyaluronic acid/chondroitin sulfate on healing of full-thickness tendon lacerations in rabbits. *J Orthop Res*, 1989. 7(5): 683-9.

147. Green, S., et al., The inhibition of flexor tendon adhesions. *Bull Hosp Jt Dis Orthop Inst*, 1986. 46(1): 16-21.
148. Menderes, A., et al., Prevention of peritendinous adhesions following flexor tendon injury with seprafilm. *Ann Plast Surg*, 2004. 53(6): 560-4.
149. Karakurum, G., et al., Seprafilm interposition for preventing adhesion formation after tenolysis. An experimental study on the chicken flexor tendons. *J Surg Res*, 2003. 113(2): 195-200.
150. St Onge, R., et al., A preliminary assessment of Na-hyaluronate injection into "no man's land" for primary flexor tendon repair. *Clin Orthop Relat Res*, 1980(146): 269-75.
151. Moro-oka, T., et al., Mixture of hyaluronic acid and phospholipid prevents adhesion formation on the injured flexor tendon in rabbits. *J Orthop Res*, 2000. 18(5): 835-40.
152. Wiig, M., S.O. Abrahamsson, and G. Lundborg, Effects of hyaluronan on cell proliferation and collagen synthesis: a study of rabbit flexor tendons in vitro. *J Hand Surg [Am]*, 1996. 21(4): 599-604.
153. Hagberg, L., Exogenous hyaluronate as an adjunct in the prevention of adhesions after flexor tendon surgery: a controlled clinical trial. *J Hand Surg [Am]*, 1992. 17(1): 132-6.
154. Taras, J.S. and M.J. Lamb, Treatment of flexor tendon injuries: surgeons' perspective. *J Hand Ther*, 1999. 12(2): 141-8.
155. Hagberg, L., A. Tengblad, and B. Gerdin, Elimination of exogenously injected sodium-hyaluronate from rabbit flexor tendon sheaths. *J Orthop Res*, 1991. 9(6): 792-7.
156. Hagberg, L. and B. Gerdin, Sodium hyaluronate as an adjunct in adhesion prevention after flexor tendon surgery in rabbits. *J Hand Surg [Am]*, 1992. 17(5): 935-41.
157. Isik, S., et al., Prevention of restrictive adhesions in primary tendon repair by HA-membrane: experimental research in chickens. *Br J Plast Surg*, 1999. 52(5): 373-9.
158. Weiss, C., et al., The role of Na-hylan in reducing postsurgical tendon adhesions. *Bull Hosp Jt Dis Orthop Inst*, 1986. 46(1): 9-15.
159. Weiss, C., et al., The role of Na-hylan in reducing postsurgical tendon adhesions: Part 2. *Bull Hosp Jt Dis Orthop Inst*, 1987. 47(1): 31-9.
160. Wang, S.J., et al., [The clinical study of adhesion prevention of sodium hyaluronate in flexor tendon surgery]. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi*, 2002. 16(1): 28-30.
161. Banes, A.J., et al., PDGF-BB, IGF-I and mechanical load stimulate DNA synthesis in avian tendon fibroblasts in vitro. *J Biomech*, 1995. 28(12): 1505-13.
162. Dowthwaite, G.P., et al., The effect of mechanical strain on hyaluronan metabolism in embryonic fibrocartilage cells. *Matrix Biol*, 1999. 18(6): 523-32.
163. Arnoczky, S.P., et al., Activation of stress-activated protein kinases (SAPK) in tendon cells following cyclic strain: the effects of strain frequency, strain magnitude, and cytosolic calcium. *J Orthop Res*, 2002. 20(5): 947-52.

164. Zeichen, J., M. van Griensven, and U. Bosch, The proliferative response of isolated human tendon fibroblasts to cyclic biaxial mechanical strain. *Am J Sports Med*, 2000. 28(6): 888-92.
165. Qin, T.W., et al., Adhesion strength of human tenocytes to extracellular matrix component-modified poly(DL-lactide-co-glycolide) substrates. *Biomaterials*, 2005. 26(33): 6635-42.
166. Koob, T.J., Biomimetic approaches to tendon repair. *Comp Biochem Physiol A Mol Integr Physiol*, 2002. 133(4): 1171-92.
167. Yamamoto, E., et al., Effects of static stress on the mechanical properties of cultured collagen fascicles from the rabbit patellar tendon. *J Biomech Eng*, 2002. 124(1): 85-93.
168. Ralphs, J.R., A.D. Waggett, and M. Benjamin, Actin stress fibres and cell-cell adhesion molecules in tendons: organisation in vivo and response to mechanical loading of tendon cells in vitro. *Matrix Biol*, 2002. 21(1): 67-74.
169. Hannafin, J.A., et al., Effect of stress deprivation and cyclic tensile loading on the material and morphologic properties of canine flexor digitorum profundus tendon: an in vitro study. *J Orthop Res*, 1995. 13(6): 907-14.
170. Koob, T.J., et al., Compression loading in vitro regulates proteoglycan synthesis by tendon fibrocartilage. *Arch Biochem Biophys*, 1992. 298(1): 303-12.
171. Slack, C., M.H. Flint, and B.M. Thompson, The effect of tensional load on isolated embryonic chick tendons in organ culture. *Connect Tissue Res*, 1984. 12(3-4): 229-47.
172. Gillard, G.C., et al., The influence of mechanical forces on the glycosaminoglycan content of the rabbit flexor digitorum profundus tendon. *Connect Tissue Res*, 1979. 7(1): 37-46.
173. Woo, S.L., et al., The effects of exercise on the biomechanical and biochemical properties of swine digital flexor tendons. *J Biomech Eng*, 1981. 103(1): 51-6.
174. Malaviya, P., et al., An in vivo model for load-modulated remodeling in the rabbit flexor tendon. *J Orthop Res*, 2000. 18(1): 116-25.
175. Buchanan, C.I. and R.L. Marsh, Effects of exercise on the biomechanical, biochemical and structural properties of tendons. *Comp Biochem Physiol A Mol Integr Physiol*, 2002. 133(4): 1101-7.
176. Packer, D.L., et al., An in vitro model of fibroblast activity and adhesion formation during flexor tendon healing. *J Hand Surg [Am]*, 1994. 19(5): 769-76.
177. Spicer, A.P. and J.A. McDonald, Characterization and molecular evolution of a vertebrate hyaluronan synthase gene family. *J Biol Chem*, 1998. 273(4): 1923-32.
178. Weigel, P.H., V.C. Hascall, and M. Tammi, Hyaluronan synthases. *J Biol Chem*, 1997. 272(22): 13997-4000.
179. Rilla, K., et al., Changed lamellipodial extension, adhesion plaques and migration in epidermal keratinocytes containing constitutively expressed sense and antisense hyaluronan synthase 2 (Has2) genes. *J Cell Sci*, 2002. 115(Pt 18): 3633-43.
180. Falk, K., et al., Prevention of adhesions by surfactants and cellulose derivatives in mice. *Eur J Surg*, 2001. 167(2): 136-41.
181. Khitrina, G.V., [The effect of hyaluronidase on adhesive processes in the pleural cavity under experimental conditions]. *Biull Eksp Biol Med*, 1968. 65(4): 115-7.

182. Stoehr, B.J., J.E. Gutierrez, and A.S. Close, Effect of intraperitoneal hyaluronidase on the reformation of intestinal adhesions. *Am J Surg*, 1966. 111(6): 881-3.
183. Adamian, L.V., O.A. Mynbaev, and V.M. Strugatskii, [An experimental validation of hyaluronidase electrophoresis for the prevention of postoperative adhesions]. *Vopr Kurortol Fizioter Lech Fiz Kult*, 1995(3): 18-20.
184. Zimmerman, E., B. Geiger, and L. Addadi, Initial stages of cell-matrix adhesion can be mediated and modulated by cell-surface hyaluronan. *Biophys J*, 2002. 82(4): 1848-57.
185. Yang, C., et al., Tendon surface modification by chemically modified HA coating after flexor digitorum profundus tendon repair. *J Biomed Mater Res B Appl Biomater*, 2004. 68(1): 15-20.
186. Kelly, M.A., R.W. Moskowitz, and J.R. Lieberman, Hyaluronan therapy: looking toward the future. *Am J Orthop*, 2004. 33(2 Suppl): 23-8.
187. Pitt, W.G., et al., Attachment of hyaluronan to metallic surfaces. *J Biomed Mater Res A*, 2004. 68(1): 95-106.
188. Bulpitt, P. and D. Aeschlimann, New strategy for chemical modification of hyaluronic acid: preparation of functionalized derivatives and their use in the formation of novel biocompatible hydrogels. *J Biomed Mater Res*, 1999. 47(2): 152-69.
189. Tsai, S.W., et al., Preparation and evaluation of a hyaluronate-collagen film for preventing post-surgical adhesion. *J Int Med Res*, 2005. 33(1): 68-76.