

Yale University

EliScholar – A Digital Platform for Scholarly Publishing at Yale

Cowles Foundation Discussion Papers

Cowles Foundation

9-1-1975

The Value of a Nonstandard Competitive Allocation

Donald J. Brown

Peter A. Loeb

Follow this and additional works at: <https://elischolar.library.yale.edu/cowles-discussion-paper-series>



Part of the [Economics Commons](#)

Recommended Citation

Brown, Donald J. and Loeb, Peter A., "The Value of a Nonstandard Competitive Allocation" (1975). *Cowles Foundation Discussion Papers*. 639.

<https://elischolar.library.yale.edu/cowles-discussion-paper-series/639>

This Discussion Paper is brought to you for free and open access by the Cowles Foundation at EliScholar – A Digital Platform for Scholarly Publishing at Yale. It has been accepted for inclusion in Cowles Foundation Discussion Papers by an authorized administrator of EliScholar – A Digital Platform for Scholarly Publishing at Yale. For more information, please contact elischolar@yale.edu.

COWLES FOUNDATION FOR RESEARCH IN ECONOMICS

AT YALE UNIVERSITY

Box 2125, Yale Station
New Haven, Connecticut 06520

COWLES FOUNDATION DISCUSSION PAPER NO. 406

Note: Cowles Foundation Discussion Papers are preliminary materials circulated to stimulate discussion and critical comment. Requests for single copies of a Paper will be filled by the Cowles Foundation within the limits of the supply. References in publications to Discussion Papers (other than mere acknowledgment by a writer that he has access to such unpublished material) should be cleared with the author to protect the tentative character of these papers.

THE VALUE OF A NONSTANDARD COMPETITIVE ALLOCATION

Donald J. Brown and Peter A. Loeb

September 30, 1975

THE VALUE OF A NONSTANDARD COMPETITIVE ALLOCATION*

by

Donald J. Brown** and Peter A. Loeb***

September, 1975

I. Introduction

In [1], Aumann gives a heuristic proof of the value equivalence theorem for exchange economies with a non-atomic continuum of traders. His nonrigorous argument is based on a naive notion of infinitesimals. We shall prove a value equivalence theorem for nonstandard exchange economies, using Robinson's calculus of infinitesimals, i.e. nonstandard calculus. Our proof can be viewed as a rigorous version of Aumann's original intuitive argument.

As a corollary to our value equivalence theorem we obtain a standard limit theorem. This theorem gives a characterization of competitive allocations in large standard exchange economies as approximate value allocations. Moreover, it extends Champsaur's results on the value of competitive allocations in replicated exchange economies [6].

Nonstandard exchange economies and the associated non-standard concepts of the core and competitive equilibrium were

*The research in this paper was supported in part by the National Science Foundation and the Ford Foundation.

**Cowles Foundation, Yale University

***Mathematics Departments, Yale University and the University of Illinois, Urbana-Champaign

first defined in Brown-Robinson [3]. In that paper, they proved the equivalence between the nonstandard core and the set of nonstandard competitive allocations, i.e. an allocation is in the core iff it is a competitive allocation. In a second paper [4], using the core equivalence theorem for nonstandard exchange economies, they showed that core allocations in large standard economies are approximate competitive allocations. The interested reader is referred to either of these two papers for an introduction to nonstandard analysis and a discussion of nonstandard exchange economies.

We should note that the existence of a value allocation follows from the value equivalence theorem, the core equivalence theorem, and the existence of nonstandard competitive equilibria shown by Brown in [5].

Before stating and proving our two major theorems, we shall need the following definitions.

II. Definitions

Let *R_d be the d -fold Cartesian product of *R , the nonstandard extension of R , and ${}^*\Omega_d$ be the positive orthant of *R_d . If \bar{x} and \bar{y} are vectors in *R_d , then we shall write $\bar{x} \simeq \bar{y}$ when the distance between x and y is infinitesimal in the metric defined by the sup norm. $\bar{x} \geq \bar{y}$ means $x_i \geq y_i$ for all i ; $\bar{x} > \bar{y}$ means $\bar{x} \geq \bar{y}$ and $x_i > y_i$ for some i ; $\bar{x} \gg \bar{y}$ means $x_i > y_i$ for all i . $\bar{x} \gtrsim \bar{y}$ means $x_i \geq y_i$ or $x_i \simeq y_i$ for all i ; $\bar{x} \gtrsim \bar{y}$ means $\bar{x} \gtrsim \bar{y}$ and x_i

is greater than y_i by a noninfinitesimal amount for some i ; $\bar{x} \gg \bar{y}$ means x_i is greater than y_i by a noninfinitesimal amount for all i . The vector which is 0 in all components except the i^{th} is denoted by \bar{e}^i .

By the norm of a vector $\bar{x} = (\bar{x}_1, \dots, \bar{x}_d)$, we mean the sup norm $\|\bar{x}\| = \max_{1 \leq j \leq d} |\bar{x}_j|$. A nonstandard vector is said to be finite or near standard if its sup norm is less than some standard number. If \bar{x} is a finite vector, then there exists a unique standard vector, called the standard part of \bar{x} , denoted by ${}^o\bar{x}$, where ${}^o\bar{x} \simeq \bar{x}$.

$*N$ will denote the nonstandard extension of N , the integers, and $*N - N$ is the set of infinite integers. If δ is an infinitesimal, then we shall often write $\delta \simeq 0$, with similar notation used for infinitesimal vectors.

Let T be an internal star-finite set. T will be called the set of traders or agents. We shall always assume that T is infinite, i.e. the internal cardinality $|T| = \omega \in *N - N$.

A coalition is an internal subset of traders.

A coalition S is negligible if $|S|/\omega \simeq 0$.

All agents are assumed to have the same consumption set which is $*\Omega_d$.

An assignment is an internal map from T into $*\Omega_d$.

A trader is defined by his initial endowment, an element of $*\Omega_d$, and his preference relation, a binary relation on $*\Omega_d$.

A nonstandard exchange economy, \mathcal{E} , is a pair $\langle I, P \rangle$ where $I(t)$ is an assignment and $P(t)$ is an internal map from T into the family of internal binary relations on ${}^*\Omega_d$. We will often denote $P(t)$ as $>_t$. That is, $>_t$ is the preference relation of trader t and $I(t)$ is his initial endowment.

An allocation, X , is an assignment such that $X(t)$ is finite for each $t \in T$. It follows that the norm $\|X(t)\|$ is uniformly bounded.

An allocation X is feasible for a coalition S if

$$\frac{1}{|S|} \sum_{t \in S} X(t) \lesssim \frac{1}{|S|} \sum_{t \in S} I(t).$$

An allocation X is strictly feasible for a coalition S if

$$\frac{1}{|S|} \sum_{t \in S} X(t) \leq \frac{1}{|S|} \sum_{t \in S} I(t), \text{ i.e. } \sum_{t \in S} X(t) \leq \sum_{t \in S} I(t).$$

If $S = T$, then we will say that X is feasible or strictly feasible.

If $\bar{x}, \bar{y} \in {}^*\Omega_d$, then $\bar{x} >_t \bar{y}$ iff for all $\bar{w} \simeq \bar{x}$ and for all $\bar{z} \simeq \bar{y}$, $\bar{w} >_t \bar{z}$. If $\bar{p} \in {}^*\Omega_d$, then $B_{\bar{p}}(t) = \{\bar{x} \in {}^*\Omega_d : \bar{p} \cdot \bar{x} \lesssim \bar{p} \cdot I(t)\}$. Given a feasible allocation $X(t)$, $\langle X, \bar{p} \rangle$ is a competitive equilibrium for the economy $\mathcal{E} = \langle I(t), \{>_t\}_{t \in T} \rangle$ iff \bar{p} is finite, $\bar{p} \gg 0$ and there exists a coalition T_0 , with $|T_0|/|T| \simeq 0$, such that for all $t \in T - T_0$, $X(t) \in B_{\bar{p}}(t)$ and $y \notin B_{\bar{p}}(t)$ for any $\bar{y} >_t X(t)$. If $\langle X, \bar{p} \rangle$ is a competitive equilibrium, then we say that X is a competitive allocation and \bar{p} is a competitive price system.

If $>_t$ is a preference relation over $*\Omega_d$, then a utility (function) for $>_t$ is an internal map $u_t : *\Omega_d \rightarrow *R$ such that for all $\bar{x}, \bar{y} \in *\Omega_d$, $\bar{x} >_t \bar{y}$ iff $u_t(\bar{x}) > u_t(\bar{y})$.

If $\{>_t\}_{t \in T}$ is an internal family of preference relations, then an internal family of utilities $\{u_t\}_{t \in T}$ is said to represent $\{>_t\}_{t \in T}$ if for each $t \in T$, u_t is a utility function for $>_t$.

If $\{u_t\}_{t \in T}$ is a representing family for $\{>_t\}_{t \in T}$, then \mathcal{E} can be expressed as $\langle I(t), \{u_t\}_{t \in T} \rangle$. We now use \mathcal{E} to define a star-finite internal game \mathcal{G} , in characteristic form, over T . Let $\mathcal{A}(T)$ be the family of internal subsets of T or coalitions of T . Then

$\mathcal{G} = \langle V, \mathcal{A}(T) \rangle$ where $V : \mathcal{A}(T) \rightarrow *R$ is defined as follows: Given $S \in \mathcal{A}(T)$, $V(S) = \max \frac{1}{\omega} \sum_{t \in S} u_t(Y(t))$ where Y ranges over all assignments with $\sum_{t \in S} Y(t) \leq \sum_{t \in S} I(t)$.

The Shapley-value is an a priori evaluation of a finite game for each player. A brief discussion of the Shapley-value and its properties is given in Appendix A of [2]. By transfer, the Shapley-value of \mathcal{G} is well defined and we denote it by ϕ_V . Of course, \mathcal{G} and hence ϕ_V depends on $\{u_t\}_{t \in T}$ and in general different representing families define different games.

A value allocation with respect to $\mathcal{E} = \langle I(t), \{u_t\}_{t \in T} \rangle$ is a feasible allocation X such that for each $S \in \mathcal{A}(T)$, $\sum_{t \in S} \phi_V(t) \simeq \frac{1}{\omega} \sum_{t \in S} u_t(X(t))$. It is important to note that a value allocation depends on the representation of the preferences $\{>_t\}_{t \in T}$.

Debreu [7] defines a (standard) smooth preference relation over Ω_d as a binary relation over Ω_d having a quasi-concave C^2 standard utility with a strictly positive gradient and indifference surfaces with everywhere positive Gaussian curvature.

A uniformly smooth family of (standard) preferences was first defined by Aumann[1]. In particular, he considered the following example of a uniformly smooth family, which we shall call a precompact smooth family. A family of standard utilities $\{u_s\}_{s \in S}$ on Ω_d (i.e. for all $s \in S$, $u_s : \Omega_d \rightarrow \mathbb{R}$) is said to be a precompact smooth family if:

- (a) Each u_s is smooth (in the sense of Debreu).
- (b) The u_s are bounded over Ω_d uniformly in s .
- (c) $\{u_s\}_{s \in S}$ is contained in a compact subset of the space \mathcal{U} of C^2 utility functions on Ω_d endowed with the topology of C^2 uniform convergence on compact sets.

If $\bar{x} \in \Omega_d$, then a C^1 utility (function) u on Ω_d is called concave over Ω_d at \bar{x} (or concave at \bar{x}) if for all $\bar{y} \in \Omega_d$, $u(\bar{y}) - u(\bar{x}) \leq \nabla u(\bar{x}) \cdot (\bar{y} - \bar{x})$ where $\nabla u(\bar{x})$ is the gradient of u evaluated at \bar{x} .

III. Assumptions

- (1) $|T| = \omega \in {}^*N - N$.
- (2) $I(t)$ is an allocation.
- (3) $\frac{1}{\omega} \sum_{t \in T} I(t) \gg \bar{0}$.
- (4) For all $t \in T$, $I(t) \not\leq \bar{0}$.
- (5) The preferences $\{>_t\}_{t \in T}$ are represented by an internal

family of utility functions $\{u_t\}_{t \in T}$ where $\{u_t\}_{t \in T}$ is contained in the nonstandard extension of a precompact smooth family of standard utility functions.

By a representing family of utilities, we shall mean a family satisfying assumption (5).

Aumann in [1], Lemma 15.1, has shown that if $\{u_s\}_{s \in S}$ is a precompact smooth family of standard utility functions, then for every $\gamma > 0$ there exists a family of utilities $\{\tilde{u}_s\}_{s \in S}$ such that:

- (a) The \tilde{u}_s are uniformly bounded.
- (b) The \tilde{u}_s are C^1 on Ω_d .
- (c) The gradients $\nabla \tilde{u}_s$ are, on compact sets, uniformly bounded and uniformly positive, i.e. for every compact set K there exists vectors \bar{a} and \bar{b} , where $\bar{a} \gg \bar{0}$ and $\bar{b} \gg \bar{0}$ such that for all $s \in S$, for all $\bar{y} \in K$, $\bar{a} \leq \nabla \tilde{u}_s(\bar{y}) \leq \bar{b}$.
- (d) Each \tilde{u}_s is concave over Ω_d at each \bar{x} such that $\|\bar{x}\| \leq \gamma$.

A family of utilities satisfying (a), (b), (c), and (d), above shall be called essentially concave in $B_\gamma(\bar{0}) = \{\bar{x} \in \Omega_d : \|\bar{x}\| \leq \gamma\}$.

By a representing family of utilities $\{u_t\}_{t \in T}$ essentially concave in ${}^*B_\gamma(\bar{0})$, we mean an internal subset of the nonstandard extension of a family of standard utilities essentially concave in $B_\gamma(\bar{0})$. Moreover, this family of

standard utilities is assumed to be compact in the topology of C^1 uniform convergence on compact sets.

IV. Statement of the Main Theorem

Let $\mathcal{E} = \langle I(t), \{>_t\}_{t \in T} \rangle$ be a nonstandard exchange economy satisfying the assumptions in Section III.

Theorem 1. (a) Given a nonstandard competitive allocation X for \mathcal{E} , there is a $\gamma > 0$ such that if $\{u_t\}_{t \in T}$ is a representing family of utilities essentially concave in ${}^*B_\gamma(\bar{O})$, then there is an internal family of weights $\{\alpha_t\}_{t \in T}$ with $\alpha_t > 0$ for each t such that X is a value allocation with respect to the family of utilities $\{\alpha_t u_t\}_{t \in T}$.

(b) There exists a finite γ such that for every representing family of utilities $\{u_t\}_{t \in T}$, for all coalitions S and every allocation $X(t)$ strictly feasible for S , if $\sum_{t \in S} u_t(Y(t)) \leq \sum_{t \in S} u_t(X(t))$ for all allocations $Y(t)$ strictly feasible for S , then $\max_{t \in S} \|X(t)\| < \gamma$.

(c) If $\{u_t\}_{t \in T}$ is a representing family of utilities, essentially concave in ${}^*B_\gamma(\bar{O})$ where γ is given in (b), and X is a value allocation with respect to $\{u_t\}_{t \in T}$ then X is a competitive allocation.

V. Proof of Main Theorem

Lemma 1. Let $\{u_t\}_{t \in T}$ be a representing family of utilities. Given a coalition $S \subset T$ and an allocation $Z(t)$, let $\mathcal{C} = \{Y: Y \text{ is an assignment and } \sum_{t \in S} Y(t) \leq \sum_{t \in S} Z(t)\}$. If $X \in \mathcal{C}$ and $\sum_{t \in S} u_t(Y(t)) \leq \sum_{t \in S} u_t(X(t))$ for all $Y \in \mathcal{C}$, then

- (i) For every $v \in {}^*N - N$, $\max_{t \in S} \|X(t)\| < v$.
- (ii) There exists $\bar{p} \in {}^*\Omega_d$ such that
- (a) \bar{p} is finite and $\bar{p} \gg \bar{0}$,
- (b) Given $t \in S$, if $X_j(t) \neq \bar{0}$, then $\bar{p}_j = \nabla_j u_t(X(t))$, and if $X_j(t) = 0$, then $\bar{p}_j \geq \nabla_j u_t(X(t))$.

Proof. Suppose for some $s \in S$ and $j \leq d$, $X_j(s) \geq v$ where $v \in {}^*N - N$. Then $|S| \in {}^*N - N$ since Z is an allocation and $X \in \mathcal{C}$. Let $A_n = \{t \in S: \|X(t)\| < n\}$.

If $|A_n|$ is finite for all $n \in N$, then there exists a $\rho \in {}^*N - N$ such that $|A_\rho|/|S|^{1/2} \simeq 0$ and $\rho/|S|^{1/2} \simeq 0$.

Let $\|\bar{x}\|_1 = \sum_{j=1}^d \bar{x}_j$ if $\bar{x} \in {}^*\Omega_d$. Since preferences are

monotonic, we have $\sum_{t \in S} X(t) = \sum_{t \in S} Z(t)$ so $\sum_{t \in S} \|X(t)\|_1 =$

$\sum_{t \in S} \|Z(t)\|_1$. But $\frac{1}{|S|} \sum_{t \in S} \|X(t)\|_1 = \frac{1}{|S|} \sum_{t \in A_\rho} \|X(t)\|_1 +$

$\frac{1}{|S|} \sum_{t \in S - A_\rho} \|X(t)\|_1 \simeq \frac{1}{|S|} \sum_{t \in S - A_\rho} \|X(t)\|_1 \geq \frac{|S - A_\rho|}{|S|} \rho \geq \frac{1}{2} \rho$.

This contradicts the finiteness of $\frac{1}{|S|} \sum_{t \in S} \|Z(t)\|_1$. Hence,

for some $n_0 \in \mathbb{N}$, $|A_{n_0}| \in {}^*\mathbb{N} - \mathbb{N}$. Let $\theta = \min(v, |A_{n_0}|)$;

choose $B \subset A_{n_0}$ with $|B| = \theta$, and define a new allocation

$$W(t) = \begin{cases} X(t) + \bar{e}_j & \text{for } t \in B \\ X(t) - \theta \bar{e}_j & \text{for } t = s \\ X(t) & \text{otherwise.} \end{cases}$$

Since the u_t are standardly bounded, $u_s(X(s))$ and $u_s(W(s))$ differ by only a finite amount. The u_t are near standard on

any compact standard set, hence they are near standard on

$\{\bar{x} \in {}^*\Omega_d : \|\bar{x}\| \leq \max_{t \in B} \|W(t)\|\}$. Therefore, $\min_{t \in B} [u_t(W(t)) -$

$u_t(X(t))] > 0$, and $\sum_{t \in S} u_t(W(t)) > \sum_{t \in S} u_t(X(t))$ which contra-

dicts the definition of $X(t)$. Thus, for every $v \in {}^*\mathbb{N} - \mathbb{N}$,

$$\max_{t \in S} \|x(t)\| < v.$$

For part (ii) of the lemma, suppose for some $j \leq d$

and $s \in S$ that $X_j(s) > 0$. Define a vector $\Delta \bar{x}$ where

$\Delta \bar{x}_i = 0$ for $i \neq j$ and $\Delta \bar{x}_j = \delta$ with $0 < \delta < X_j(s)$. If

$r \in S$ and $r \neq s$, then the assignment $Y(t)$ where $Y(t) =$

$X(t)$ for $t \neq s$ or r ; $Y(s) = X(s) - \Delta \bar{x}$; and $Y(r) = X(r)$

+ $\Delta \bar{x}$ belongs to \mathcal{C} . Therefore, $\sum_{t \in S} u_t(Y(t)) \leq \sum_{t \in S} u_t(X(t))$.

$$\text{Thus, } \frac{u_r(X(r) + \Delta \bar{x}) - u_r(X(r))}{\delta} \leq \frac{u_s(X(s)) - u_s(X(s) - \Delta \bar{x})}{\delta},$$

which implies that $\nabla_j u_s(X(s)) \geq \nabla_j u_r(X(r))$. If $X_j(t) = 0$

for all $t \in S$, let $\bar{p}_j = \max_{t \in S} \nabla_j u_t(X(t))$. If there is a $t \in S$ with $X_j(t) > 0$, let $\bar{p}_j = \nabla_j u_t(X(t))$. Clearly, \bar{p} is well defined and $\bar{p}_j \geq \nabla_j u_t(X(t)) \geq 0$ for all $j \leq d$ and for all $t \in S$.

By Part (i), $X(t)$ is finite for all $t \in S$, thus, for all $j \leq d$ and $s \in S$, $\nabla_j u_s(X(s)) = \nabla_j^0 u_s({}^0X(s)) > 0$. Hence, for each $j \leq d$, $\bar{p}_j > 0$ and $\max_{s \in S} \nabla_j u_s(X(s))$ is finite, i.e. \bar{p} is finite.

Lemma 2. There exists a finite integer γ such that for every representing family of utilities $\{u_t\}_{t \in T}$ and every coalition S , if $X(t)$ is strictly feasible for S and $\sum_{t \in S} u_t(Y(t)) \leq \sum_{t \in S} u_t(X(t))$ for all $Y(t)$ which are strictly feasible for S , then $\max_{t \in S} \|X(t)\| < \gamma$.

Proof. Consider the internal set of integers A , where $n \in A$ if there exists a coalition S , a representing family of utilities $\{u_t\}_{t \in T}$, and allocation $X(t)$ strictly feasible for S which maximizes $\sum_{t \in S} u_t(Y(t))$ over all allocations $Y(t)$ strictly feasible for S , such that $\max_{t \in S} \|X(t)\| \geq n$. By Part (i) of Lemma 1, A contains no infinite integers. Hence, A contains only a finite number of elements. so $\gamma = 1 + \max_{j \in A} j$ gives the desired bound.

Proposition 1. Let $\{u_t\}_{t \in T}$ be a representing family of utilities, essentially concave in ${}^*B_\gamma(\bar{0})$, where γ is given in Lemma 2. Given a coalition $S \subset T$ and an allocation $Z(t)$, let $\mathcal{C} = \{Y : Y \text{ is an assignment and } \sum_{t \in S} Y(t) \leq \sum_{t \in S} Z(t)\}$, and fix $X \in \mathcal{C}$ so that $\sum_{t \in S} u_t(Y(t)) \leq \sum_{t \in S} u_t(X(t))$ for all $Y \in \mathcal{C}$. Let \bar{p} be the finite vector in ${}^*\Omega_d$ such that $\bar{p}_j = \max_{t \in S} \nabla_j u_t(X(t))$ if $X_j(t) = 0$ for all $t \in S$ and $\bar{p}_j = \nabla_j u_s(X(s))$ for each $s \in S$ such that $X_j(s) \neq 0$. Then for all $\bar{y} \in {}^*\Omega_d$, $u_t(\bar{y}) - u_t(X(t)) \leq \bar{p} \cdot (\bar{y} - X(t))$.

Proof. By Lemma 2, $\|X(t)\| < \gamma$ for all $t \in S$. Thus, by the essential concavity of the $\{u_t\}_{t \in T}$, $u_t(\bar{y}) - u_t(X(t)) \leq \nabla u(X(t)) \cdot (\bar{y} - X(t))$ for all $\bar{y} \in {}^*\Omega_d$. By Lemma 1, if $\bar{p}_j \neq \nabla_j u_t(X(t))$ then $\bar{p}_j > \nabla_j u_t(X(t))$ and $\bar{y}_j - X_j(t) \geq 0$ since $X_j(t) = 0$. Therefore, $u_t(\bar{y}) - u_t(X(t)) \leq \bar{p} \cdot (\bar{y} - X(t))$.

Definition. Given $Z, X, \{u_t\}_{t \in T}$, and \bar{p} as in Proposition 1, we write $(X, \bar{p}) \in \mathcal{M}(S, Z, \{u_t\}_{t \in T})$, and we say that X maximizes utility for S .

Lemma 3. Given $S \subset T$ where $|S| \in {}^*N - N$, let $(Y, \bar{p}) \in \mathcal{M}(S, I, \{u_t\}_{t \in T})$. If X is a feasible allocation for S such that $\frac{1}{|S|} \sum_{t \in S} u_t(X(t)) \simeq \frac{1}{|S|} \sum_{t \in S} u_t(Y(t))$, then there

exists an $S_0 \subset S$ with $|S_0|/|S| \simeq 0$ so that for all $t \in S - S_0$, $u_t(X(t)) \simeq u_t(Y(t)) + \bar{p} \cdot (X(t) - Y(t))$.

Proof. Since X and Y are feasible on S ,

$$\bar{p} \cdot \frac{1}{|S|} \sum_{t \in S} (X(t) - Y(t)) \simeq 0. \text{ Let } S_n = \{t \in S : u_t(X(t)) -$$

$$u_t(Y(t)) \leq \bar{p} \cdot (X(t) - Y(t)) - 1/n\}. \text{ Clearly, } S_n \subset S_{n+1}$$

for each $n \in \mathbb{N}$. If $n \in \mathbb{N}$ and $|S_n|/|S| \geq 1/n$, then

$$\frac{1}{|S|} \sum_{t \in S} (u_t(X(t)) - u_t(Y(t))) \leq \frac{1}{|S|} \sum_{t \in S_n} (\bar{p} \cdot (X(t) - Y(t)) - 1/n) +$$

$$\frac{1}{|S|} \sum_{t \in S - S_n} \bar{p} \cdot (X(t) - Y(t)) \leq \bar{p} \cdot \left(\frac{1}{|S|} \sum_{t \in S} (X(t) - Y(t)) \right) -$$

$$\frac{|S_n|}{|S|} \cdot \frac{1}{n} \leq -\frac{1}{n^2}.$$

Hence,
$$\frac{1}{|S|} \sum_{t \in S} u_t(X(t)) \leq \frac{1}{|S|} \sum_{t \in S} u_t(Y(t)) - 1/n^2.$$

But this is impossible. Therefore, there is a $v \in \mathbb{N} - \mathbb{N}$ with $|S_v|/|S| < 1/v$. It follows from Proposition 1 that for all $t \in S - S_v$, $u_t(X(t)) - u_t(Y(t)) \simeq \bar{p} \cdot (X(t) - Y(t))$.

Lemma 4. Let S be a subset of T with $|S| \in \mathbb{N} - \mathbb{N}$, and let X be an allocation which is feasible on S such that $\frac{1}{|S|} \sum_{t \in S} X(t) \gg \bar{0}$. Then there is an allocation $Z(t)$ which is strictly feasible on S such that $Z(t) \simeq X(t)$ for all $t \in S$.

Proof. Let $a_j = \frac{1}{|S|} \sum_{t \in S} X_j(t)$ and let $A_j = \{t \in S: X_j(t) \geq a_j/2\}$. Then $|A_j|/|S| \neq 0$. For each $t \in S$ and $j \leq d$, let

$$Z_j(t) = \begin{cases} X_j(t) + \frac{1}{|A_j|} \sum_{s \in S} (I_j(s) - X_j(s)) & \text{if } t \in A_j \\ X_j(t) & \text{otherwise.} \end{cases}$$

Since $|S|/|A_j|$ is finite and $\frac{1}{|S|} \sum_{t \in S} (I_j(t) - X_j(t)) \simeq 0$,

$Z_j(t) \simeq X_j(t)$ for all $j \leq d$ and all $t \in S$. Clearly, $Z(t)$ is strictly feasible on S .

Lemma 5. Let $(X, \bar{p}) \in \mathcal{M}(T, I, \{u_t\}_{t \in T})$ and let S be a subset of T such that $|S| \in {}^*N - N$, X is feasible on S , and

$\frac{1}{|S|} \sum_{t \in S} X(t) \gg \bar{0}$. Let $(Y, \bar{q}) \in \mathcal{M}(S, I, \{u_t\}_{t \in T})$. Then $\bar{p} \simeq \bar{q}$.

Proof. By Lemma 4, there is an allocation $Z(t)$ which is strictly feasible on S such that $Z(t) \simeq X(t)$, for all

$t \in S$. Therefore, $\frac{1}{|S|} \sum_{t \in S} u_t(X(t)) \simeq \frac{1}{|S|} \sum_{t \in S} u_t(Z(t)) \leq$

$\frac{1}{|S|} \sum_{t \in S} u_t(Y(t))$. On the other hand, $(X, \bar{p}) \in \mathcal{M}(S, X, \{u_t\}_{t \in T})$

and $\frac{1}{|S|} \sum_{t \in S} Y(t) = \frac{1}{|S|} \sum_{t \in S} I(t) \simeq \frac{1}{|S|} \sum_{t \in S} X(t)$. Again by Lemma 4

(using X instead of I) there is an allocation W with

$\sum_{t \in S} W(t) = \sum_{t \in S} X(t)$ and $W(t) \simeq Y(t)$ for each $t \in S$. There-

fore, $\frac{1}{|S|} \sum_{t \in S} u_t(Y(t)) \simeq \frac{1}{|S|} \sum_{t \in S} u_t(W(t)) < \frac{1}{|S|} \sum_{t \in S} u_t(X(t))$.

Thus, $\frac{1}{|S|} \sum_{t \in S} u_t(X(t)) \simeq \frac{1}{|S|} \sum_{t \in S} u_t(Y(t))$, so by Lemma 3 there

exists $S' \subset S$ with $\frac{|S'|}{|S|} \simeq 0$, such that for all $t \in S - S'$,

$u_t(X(t)) - u_t(Y(t)) \simeq \bar{q} \cdot (X(t) - Y(t))$. Applying

the same lemma to $Y(t)$ and (X, \bar{p}) we see that for some $S'' \subset S$ with $\frac{|S''|}{|S|} \simeq 0$ we have for all $t \in S - S''$,

$u_t(Y(t)) - u_t(X(t)) \simeq \bar{p} \cdot (Y(t) - X(t))$. Let $S_0 = S' \cup S''$,

then for all $t \in S - S_0$, $u_t(X(t)) - u_t(Y(t)) \simeq \bar{q} \cdot (X(t) - Y(t)) \simeq \bar{p} \cdot (X(t) - Y(t))$ and $|S_0|/|S| \simeq 0$.

For all $t \in T$ and for all $\bar{x} \in {}^* \Omega_d$, $u_t(\bar{x}) - u_t(X(t)) \leq \bar{p} \cdot (\bar{x} - X(t))$. Hence for all $t \in S - S_0$,

$$\begin{aligned} u_t(\bar{x}) - u_t(Y(t)) &= u_t(\bar{x}) - u_t(X(t)) + u_t(X(t)) - u_t(Y(t)) \\ &\leq \bar{p} \cdot (\bar{x} - X(t)) + \bar{p} \cdot (X(t) - Y(t)) \\ &\leq \bar{p} \cdot (\bar{x} - Y(t)). \end{aligned}$$

Fix j with $1 \leq j \leq d$. Since $\frac{1}{|S|} \sum_{t \in S} Y(t) \gg \bar{0}$ and

$\frac{|S_0|}{|S|} \simeq 0$, there exists a $t_1 \in S - S_0$ with $Y_j(t_1) \not\approx 0$.

Thus, by Part (ii) of Lemma 1, $\nabla_j u_{t_1}(Y(t_1)) = \bar{q}_j$.

On the other hand, ${}^{\circ}u_{t_1}({}^{\circ}Y(t_1) + \delta \bar{e}_j) - {}^{\circ}u_{t_1}({}^{\circ}Y(t_1)) \leq$
 ${}^{\circ}\bar{p}_j \delta$ for any real δ with $|\delta| < |{}^{\circ}Y_j(t_1)|$. Thus,
 $\bar{p}_j \simeq \nabla_j {}^{\circ}u_{t_1}({}^{\circ}Y(t_1)) \simeq \bar{q}_j$ for each $j \leq d$.

Lemma 6. Given a family of utilities $\{u_t\}_{t \in T}$ as in

Proposition 1, let V be the corresponding game and let
 $(X, \bar{p}) \in \mathcal{M}(T, I, \{u_t\}_{t \in T})$. Let S be a subset of T such
 that $|S| \in {}^*N - N$, X is feasible on S and $\frac{1}{|S|} \sum_{t \in S} X(t) > \bar{0}$.
 Then for every $t_0 \in S$, $V(S) - V(S - \{t_0\}) = \frac{1}{\omega} [u_{t_0}(X(t_0)) +$
 $\bar{p} \cdot (I(t_0) - X(t_0)) + \epsilon]$ where $\epsilon \simeq 0$.

Proof. Let $(Y, \bar{q}) \in \mathcal{M}(S, I, \{u_t\}_{t \in T})$ and let $(Z, \bar{l}) \in$
 $\mathcal{M}(S - \{t_0\}, I, \{u_t\}_{t \in T})$. By Lemma 5, $\bar{q} \simeq \bar{p} \simeq \bar{l}$. Moreover,

$$Y(t_0) + \sum_{t \in S - \{t_0\}} Y(t) = I(t_0) + \sum_{t \in S - \{t_0\}} I(t) \text{ and therefore}$$

$$\sum_{t \in S - \{t_0\}} (Y(t) - Z(t)) = \sum_{t \in S - \{t_0\}} (Y(t) - I(t)) = I(t_0) - Y(t_0).$$

By Proposition 1, $\sum_{t \in S - \{t_0\}} u_t(Y(t)) - u_t(Z(t)) \leq \bar{l} \cdot \sum_{t \in S - \{t_0\}} (Y(t) -$

$$Z(t)) = \bar{l} \cdot (I(t_0) - Y(t_0)) \simeq \bar{q} \cdot (I(t_0) - Y(t_0)) =$$

$$\bar{q} \cdot \sum_{t \in S - \{t_0\}} (Y(t) - Z(t)) \leq \sum_{t \in S - \{t_0\}} u_t(Y(t)) - u_t(Z(t)).$$

Hence, $\sum_{t \in S} u_t(Y(t)) - \sum_{t \in S - \{t_0\}} u_t(Z(t)) = u_{t_0}(Y(t_0)) +$

$$\sum_{t \in S - \{t_0\}} u_t(Y(t)) - u_t(Z(t)) \simeq u_{t_0}(Y(t_0)) + \bar{p} \cdot (I(t_0) - Y(t_0)).$$

But $u_{t_0}(Y(t_0)) \simeq u_{t_0}(X(t_0)) + \bar{p} \cdot (Y(t_0) - X(t_0))$. This follows from the fact that $\bar{p} \simeq \bar{q}$ and Proposition 1.

Hence $\omega(V(S) - V(S - \{t_0\})) = \sum_{t \in S} u_t(Y(t)) - \sum_{t \in S - \{t_0\}} u_t(Z(t)) \simeq u_{t_0}(X(t_0)) + \bar{p} \cdot (I(t_0) - X(t_0))$.

Lemma 7. For any $S \subset T$ and any $t_0 \in S$, $\omega(V(S) - V(S - \{t_0\}))$ is finite.

Proof. Let $(Y, \bar{q}) \in \mathcal{M}(S, I, \{u_t\}_{t \in T})$ and $(Z, \bar{l}) \in \mathcal{M}(S - \{t_0\}, I, \{u_t\}_{t \in T})$. As in the proof of the previous lemma we have $\sum_{t \in S - \{t_0\}} Y(t) - Z(t) = I(t_0) - Y(t_0)$. Therefore,

$$\omega(V(S) - V(S - \{t_0\})) = \sum_{t \in S} u_t(Y(t)) - \sum_{t \in S - \{t_0\}} u_t(Z(t)) =$$

$$u_{t_0}(Y(t_0)) + \sum_{t \in S - \{t_0\}} u_t(Y(t)) - u_t(Z(t)) \leq u_{t_0}(Y(t_0)) +$$

$\bar{l} \cdot (I(t_0) - Y(t_0))$ by Proposition 1.

Note that Lemma 7 implies the $\max_{\substack{S \subset T \\ t_0 \in S}} [\omega(V(S) - V(S - \{t_0\}))]$

is finite.

Lemma 8. Let $(X, p) \in \mathcal{M}(T, I, \{u_t\}_{t \in T})$ and let $n \in {}^*N - N$.

Then almost all coalitions $S \subset T$ with $|S| = n$ have the property that $\frac{1}{n} \sum_{t \in S} X(t) \approx \frac{1}{n} \sum_{t \in S} I(t) \gg \bar{0}$. (Here "almost

all" means that if \mathcal{S} is the set of coalitions of size n and \mathcal{S}_0 is the subset for which the theorem is false, then $|\mathcal{S}_0|/|\mathcal{S}| \approx 0$.)

Proof. Let $\delta = n^{-\frac{1}{4}}$. For each $k \in N$, $k^{2d} \delta \approx 0$, where d is the dimension of the commodity space. Fix $\gamma \in {}^*N - N$ so that $\gamma^{2d} \delta \approx 0$. Let $L = \max_{t \in T} (\|X(t)\| + \|I(t)\| + 1)$, and let B be the box defined by

$$B = \{\bar{x} \in {}^*\Omega_d : 0 \leq \bar{x}_j < L \quad \forall j \leq d\}.$$

Let \mathcal{B} be the set of infinitesimal boxes of the form

$$\{\bar{x} \in B : \frac{k_j}{\gamma} L \leq \bar{x}_j < \frac{k_{j+1}}{\gamma} L\},$$

where for each $j \leq d$, $0 \leq k_j \leq \gamma - 1$, and $k_j \in {}^*N$.

Divide T into equivalence classes called types:

t_1 and t_2 are of the same type if $X(t_1)$ and $X(t_2)$ are in the same box in \mathcal{B} and if $I(t_1)$ and $I(t_2)$ are in the same box in \mathcal{B} . Let m be the number of types in T ; clearly $m \leq \gamma^{2d}$. Order the types and let r_i , $1 \leq i \leq m$, be the ratio of the number of elements of the i^{th} type to $|T| = \omega$.

By Chebyshev's inequality, the probability is at most $\frac{1}{4n\delta^2}$ of obtaining an internal random sample S of size n from T for which the ratio s_i of the number of elements of type i in S to n differs from r_i by at least δ . That is,

$$P(|s_i - r_i| \geq \delta) \leq \frac{1}{4n\delta^2} = \frac{\delta^2}{4},$$

since $\delta \equiv n^{-\frac{1}{4}}$. Let $\mathcal{S} = \{S \subset T: |S| = n\}$. We have shown that there is a set $\mathcal{S}_0 \subset \mathcal{S}$ with $|\mathcal{S}_0|/|\mathcal{S}| \leq \frac{\delta^2}{4} \gamma^{2d} \approx 0$ so that for each $S \in \mathcal{S} - \mathcal{S}_0$, $\max_i |s_i - r_i| < \delta$.

Choose one representative $t_i \in T$ for each type. Using the ordering on T , let t_{ij} be the j^{th} element in the i^{th} type; let $\bar{\alpha}_{ij} = X(t_{ij}) - X(t_i)$ and $\bar{\beta}_{ij} = I(t_{ij}) - I(t_i)$. Clearly, $\|\bar{\alpha}_{ij}\| \approx 0$ and $\|\bar{\beta}_{ij}\| \approx 0$. Given $S \in \mathcal{S}$, and s_i , the ratio of the number of elements in S of type i to n , let j_i run through the elements of type i in S when $s_i \neq 0$. Then

$$\begin{aligned} & \frac{1}{n} \sum_{t \in S} X(t) - I(t) \\ &= \frac{1}{n} \sum_{i: s_i \neq 0} \sum_{j_i=1}^{ns_i} [(X(t_i) + \bar{\alpha}_{ij_i}) - (I(t_i) + \bar{\beta}_{ij_i})] \\ &= \sum_{i=1}^m s_i (X(t_i) - I(t_i)) + \frac{1}{n} \sum_{i: s_i \neq 0} \sum_{j_i=1}^{ns_i} (\bar{\alpha}_{ij_i} - \bar{\beta}_{ij_i}). \end{aligned}$$

Since n is at least the number of terms in the second sum on the right side of the last equation and the average of infinitesimal vectors is infinitesimal, we have

$$\frac{1}{n} \sum_{t \in S} X(t) - I(t) \simeq \sum_{i=1}^m s_i (X(t_i) - I(t_i)).$$

By similar reasoning, we have

$$\bar{O} = \frac{1}{\omega} \sum_{t \in T} X(t) - I(t) \simeq \sum_{i=1}^m r_i (X(t_i) - I(t_i)).$$

Now if $S \in \mathcal{I} - \mathcal{I}_0$, then

$$\begin{aligned} \left\| \frac{1}{n} \sum_{t \in S} X(t) - I(t) \right\| &\simeq \left\| \sum_{i=1}^m (s_i + (s_i - r_i)) (X(t_i) - I(t_i)) \right\| \\ &\leq \max_i |s_i - r_i| \cdot \max_i \|X(t_i) - I(t_i)\| \cdot m \\ &\leq \delta \gamma^{2d} \cdot \max_i \|X(t_i) - I(t_i)\| \simeq 0. \end{aligned}$$

On the other hand,

$$\begin{aligned} \frac{1}{\omega} \sum_{t \in T} I(t) &= \frac{1}{\omega} \sum_{i=1}^m \sum_{j_i=1}^{\omega r_i} (I(t_i) + \bar{\beta}_{ij_i}) \\ &\simeq \sum_{i=1}^m r_i I(t_i) \gg \bar{O}, \end{aligned}$$

and similarly,

$$\frac{1}{n} \sum_{t \in S} I(t) \sim \sum_{i=1}^m s_i I(t_i).$$

Since

$$\left\| \sum_{i=1}^m (s_i - r_i) I(t_i) \right\| \leq \delta \gamma^{2d} \cdot \max_i \| I(t_i) \| \sim 0,$$

$$\frac{1}{n} \sum_{t \in S} I(t) \gg \bar{0}.$$

Proposition 2. Given a family of utilities $\{u_t\}_{t \in T}$ as in Proposition 1 and the corresponding game V , if $(X, \bar{p}) \in \mathcal{M}(T, I, \{u_t\}_{t \in T})$ then for all $t_0 \in T$ and for almost all $S \subset T$ we have $\omega(V(S) - V(S - \{t_0\})) \sim u_{t_0}(X(t_0)) +$

$\bar{p} \cdot (I(t_0) - X(t_0))$. Thus, $\phi_V(t_0) = E[V(S) - V(S - \{t_0\})] =$

$\frac{1}{\omega} [u_{t_0}(X(t_0)) + \bar{p} \cdot (I(t_0) - X(t_0)) + \epsilon]$, where $\epsilon \sim 0$.

(Here E is the expectation operator.)

Proof. The formula for the Shapley-value is given by transfer as

$$\phi_V(t) = \frac{1}{\omega} \sum_{j=1}^{\omega-1} \frac{1}{\binom{\omega-1}{j}} \left(\sum_{\substack{R \subset T - \{t_0\} \\ |R|=j}} [V(R \cup \{t_0\}) - V(R)] \right).$$

Fix $k \in {}^*N - N$ such that $k/\omega \sim 0$. Since by Lemma 7

$V(R \cup \{t_0\}) - V(R)$ has a finite upper bound, we have $\phi_V(t)$

$$\approx \frac{1}{\omega} \sum_{j=k}^{\omega-1} \left(\frac{1}{\binom{\omega-1}{j}} \right) \left(\sum_{\substack{R \subset T - \{t_0\} \\ |R|=j}} [V(R \cup \{t_0\}) - V(R)] \right). \text{ For infinite}$$

coalitions R , e.g. $|R| \geq k$, we can apply Lemmas 6, 7, and 8 and obtain for fixed $j \geq k$

$$\begin{aligned} & \left(\frac{1}{\binom{\omega-1}{j}} \right) \sum_{\substack{R \subset T - \{t_0\} \\ |R|=j}} [V(R \cup \{t_0\}) - V(R)] \approx \\ & \frac{1}{\omega} [u_{t_0}(X(t_0)) + \bar{p} \cdot (I(t_0) - X(t_0))]. \end{aligned}$$

Therefore

$$\begin{aligned} \phi_V(t_0) & \approx \frac{\omega-k}{\omega^2} [u_{t_0}(X(t_0)) + \bar{p} \cdot (I(t_0) - X(t_0))] \\ & \approx \frac{1}{\omega} [u_{t_0}(X(t_0)) + \bar{p} \cdot (I(t_0) - X(t_0))]. \end{aligned}$$

We now prove Part c of Theorem 1: If Y is a value allocation with respect to an appropriate family of utilities, then Y is a competitive allocation.

If Y is a value allocation with respect to a family of utilities as in Proposition 1, then by definition

$$\frac{1}{\omega} \sum_{t \in T} u_t(Y(t)) \approx \sum_{t \in T} \phi_V(t) = V(T) = \frac{1}{\omega} \sum_{t \in T} u_t(X(t))$$

where $(X, \bar{p}) \in \mathcal{M}(T, I, \{u_t\}_{t \in T})$. Thus there exists by Lemma 3 a set $T_0 \subset T$ with $|T_0|/|T| \approx 0$ such that for

all $t \in T - T_0$, $u_t(Y(t)) \simeq u_t(X(t)) + \bar{p} \cdot (Y(t) - X(t))$.

Hence by Proposition 2,

$$\begin{aligned} u_t(Y(t)) &\simeq u_t(X(t)) + \bar{p}(I(t) - X(t)) + \bar{p}(Y(t) - I(t)) \\ &\simeq \omega[\phi_V(t)] + \bar{p} \cdot (Y(t) - I(t)). \end{aligned}$$

Therefore,

$$\phi_V(t) \simeq \frac{1}{\omega} [u_t(Y(t)) + \bar{p}(I(t) - Y(t))]$$

for all $t \in T - T_0$.

We claim that $\bar{p} \cdot Y(t) \simeq \bar{p} \cdot I(t)$ for almost all $t \in T$. Let

$$S_n = \{t \in T - T_0 : \bar{p} \cdot Y(t) \geq \bar{p} \cdot I(t) + 1/n\}$$

and

$$V_n = \{t \in T - T_0 : \bar{p} \cdot Y(t) \leq \bar{p} \cdot I(t) - 1/n\}$$

Then $S_n \subset S_{n+1}$ and $V_n \subset V_{n+1}$ for each $n \in \mathbb{N}$. Assume

that $|S_n|/|T-T_0| \geq 1/n$, then by definition, $\frac{1}{\omega} \sum_{t \in S_n} u_t(Y(t)) \simeq$

$$\sum_{t \in S_n} \phi_V(t) = \frac{1}{\omega} \sum_{t \in S_n} (u_t(Y(t)) + \bar{p} \cdot (I(t) - Y(t)) + \epsilon_t),$$

where $\epsilon_t \simeq 0$. Thus, $\frac{1}{\omega} \sum_{t \in S_n} \bar{p} \cdot (I(t) - Y(t)) \simeq 0$. But $\bar{p} \cdot I(t) -$

$\bar{p} \cdot Y(t) \leq -\frac{1}{n}$ for all $t \in S_n$, so $\frac{1}{\omega} \sum_{t \in S_n} \bar{p} \cdot (I(t) -$

$Y(t)) \leq \frac{|S_n|}{\omega} (-1/n) \leq -1/n^2$, which is impossible if $n \in \mathbb{N}$.

Therefore, for some $v \in {}^*N - N$. $|S_v|/|T-T_0| < 1/v$ and similarly $|U_v|/|T-T_0| < 1/v$. That is, $\bar{p} \cdot Y(t) \simeq \bar{p} \cdot I(t)$ except for $t \in S_v \cup U_v \cup T_0$. Thus, given $t \in T - [S_v \cup U_v \cup T_0]$ and $\bar{y} \in {}^*\Omega_d$ with $\bar{p} \cdot \bar{y} \lesssim \bar{p} \cdot I(t)$, we have $\bar{p} \cdot \bar{y} \lesssim \bar{p} \cdot Y(t)$, whence $u_t(\bar{y}) - u_t(Y(t)) = [u_t(\bar{y}) - u_t(X(t))] - [u_t(Y(t)) - u_t(X(t))] \lesssim \bar{p} \cdot (\bar{y} - X(t)) - \bar{p} \cdot (Y(t) - X(t)) = \bar{p} \cdot (\bar{y} - Y(t)) \lesssim 0$. That is, $u_t(\bar{y}) \lesssim u_t(Y(t))$.

Part b of Theorem 1 is given by Lemma 2.

We now prove Part a of Theorem 1.

Let (Y, \bar{p}) be a competitive equilibrium. Let $\{\tilde{u}_t\}_{t \in T}$ be a representing family of utilities essentially concave in ${}^*B_\gamma(\bar{0})$ where if $\gamma_1 = [\max_{t \in T} \bar{p} \cdot I(t) / \min_{1 \leq j \leq d} \bar{p}_j]$ and γ_2 is the constant in Lemma 2, then $\gamma = 1 + \gamma_1 + \gamma_2$. Let $X(t)$ be a point maximizing $\tilde{u}_t(\bar{y})$ on the budget hyperplane $B_{\bar{p}}(t) = \{\bar{y} \in {}^*\Omega_d : \bar{p} \cdot \bar{y} = \bar{p} \cdot I(t)\} \subset {}^*B_\gamma(\bar{0})$. Since (Y, \bar{p}) is a competitive equilibrium, there exists $T_0 \subset T$ such that for all $t \in T - T_0$, $\tilde{u}_t(Y(t)) \simeq \tilde{u}_t(X(t))$ and $|T_0|/|T| \simeq 0$.

Given $t \in T$, let j_0 be the first $j \leq d$ such that $X_{j_0}(t) \geq X_j(t)$, and let $\alpha_t = \bar{p}_{j_0} / \nabla_{j_0} \tilde{u}_t(X(t))$. Then $\alpha_t \not\approx 0$ is finite since $\bar{p} \not\approx \bar{0}$ and $\nabla_{j_0} \tilde{u}_t(\bar{x}) \not\approx 0$ and $\nabla_{j_0} \tilde{u}_t(\bar{x})$ is finite when $\bar{x} \in B_{\bar{p}}(t)$. Let

$u_t = \alpha_t \tilde{u}_t$; then $\nabla_{j_0} u_t(X(t)) = \bar{p}_{j_0}$. Let $\bar{x} = X(t)$, and

for $j \neq j_0$, fix $\bar{y} \in {}^* \Omega_d$ with $\bar{p} \cdot \bar{y} = \bar{p} \cdot I$ and $\bar{y}_i =$

\bar{x}_i if $i \neq j$ and $i \neq j_0$. Then $\nabla u_t(\bar{x}) \cdot (\bar{y} - \bar{x}) =$

$\bar{p}_{j_0}(\bar{y}_{j_0} - \bar{x}_{j_0}) + \nabla_j u_t(\bar{x}) \cdot (\bar{y}_j - \bar{x}_j) \leq 0$, since the directional

derivative at $X(t)$ along the budget plane is ≤ 0 . But

$\bar{p}_{j_0}(\bar{y}_{j_0} - \bar{x}_{j_0}) = -\bar{p}_j(\bar{y}_j - \bar{x}_j)$, so $\nabla_j u_t(\bar{x}) \cdot (\bar{y}_j - \bar{x}_j) -$

$\bar{p}_j(\bar{y}_j - \bar{x}_j) \leq 0$. If $X_j(t) > 0$, then $\bar{p}_j = \nabla_j u_t(\bar{x})$. Other-

wise we have $\nabla_j u_t(\bar{x}) \leq \bar{p}_j$.

We now have for any $\bar{y} \in {}^* \Omega_d$, $u_t(\bar{y}) - u_t(X(t)) \leq$

$\bar{p} \cdot (\bar{y} - X(t)) = \bar{p} \cdot \bar{y} - \bar{p} \cdot I(t)$, for all $t \in T$.

Let $(Z, \bar{q}) \in \mathcal{M}(T, I, \{u_t\}_{t \in T})$. Then

$\sum_{t \in T} u_t(Z(t)) - u_t(X(t)) \leq \sum_{t \in T} \bar{p} \cdot (Z(t) - I(t)) =$

$\bar{p} \cdot \sum_{t \in T} Z(t) - I(t) = 0$, and so $\frac{1}{\omega} \sum_{t \in T} u_t(Z(t)) \leq \frac{1}{\omega} \sum_{t \in T} u_t(X(t))$

$\approx \frac{1}{\omega} \sum_{t \in T} u_t(Y(t))$. On the other hand, $\frac{1}{\omega} \sum_{t \in T} u_t(Y(t)) -$

$u_t(Z(t)) \leq \frac{1}{\omega} \sum_{t \in T} \bar{q} \cdot (Y(t) - Z(t)) = \bar{q} \cdot [\frac{1}{\omega} \sum_{t \in T} Y(t) - Z(t)] =$

$\bar{q} \cdot [\frac{1}{\omega} \sum_{t \in T} Y(t) - I(t)] \approx 0$. Hence $\frac{1}{\omega} \sum_{t \in T} u_t(Z(t))$

$\approx \frac{1}{\omega} \sum_{t \in T} u_t(Y(t))$ which implies that $\frac{1}{\omega} \sum_{t \in T} u_t(Z(t))$

$\approx \frac{1}{\omega} \sum_{t \in T} u_t(X(t))$. Since $\bar{p} \cdot (\frac{1}{|T|} \sum_{t \in T} (X(t) - Z(t))) =$

$\bar{p} \cdot \left(\frac{1}{|T|} \sum_{t \in T} (X(t) - I(t)) \right) = 0$, we can use the proof

of Lemma 3 to show that for almost all $t \in T$, $u_t(Z(t)) - u_t(X(t)) \simeq \bar{p} \cdot (Z(t) - X(t))$. But for all $t \in T$ and all $\bar{x} \in {}^* \Omega_d$, $u_t(\bar{x}) - u_t(X(t)) \leq \bar{p} \cdot (\bar{x} - X(t))$; hence $u_t(\bar{x}) - u_t(Z(t)) \leq \bar{p} \cdot (\bar{x} - Z(t))$ for almost all $t \in T$. Using the argument at the end of the proof of Lemma 5, we see that $\bar{p} \simeq \bar{q}$.

By Proposition 2, if V is the game associated with $\{u_t\}_{t \in T}$ then for all $t \in T$, $\omega[\phi_V(t)] \simeq u_t(Z(t)) + \bar{q} \cdot (I(t) - Z(t))$. But for almost all $t \in T$, $u_t(Z(t)) + \bar{q} \cdot (I(t) - Z(t)) \simeq u_t(Z(t)) + \bar{p} \cdot (X(t) - Z(t)) + \bar{p} \cdot (I(t) - X(t)) \simeq u_t(X(t))$. Therefore, $\omega[\phi_V(t)] = u_t(Y(t)) + \epsilon_t$ where $\epsilon_t \simeq 0$ for almost all $t \in T$ and ϵ_t is finite for all $t \in T$ by Proposition 2.

Hence for all $S \subset T$, $\sum_{t \in S} \phi_V(t) \simeq \frac{1}{\omega} \sum_{t \in S} u_t(Y(t))$.

That is, $Y(t)$ is a value allocation.

This completes the proof of the main theorem, Theorem 1.

VI. A Limit Theorem

A standard exchange economy \mathcal{E} of size $m \in \mathbb{N}$ consists of m traders, whose initial endowments and preferences are restricted to the standard commodity space

Ω_d ; here, Ω_d is the d -fold Cartesian product of R . Let $\mathcal{E} = \langle I(t), >_t \rangle$ where for all t , $I(t) \in \Omega_d$ and $>_t$ is a binary relation on Ω_d . The notation $t \in \mathcal{E}$ will refer to the t^{th} trader's endowment, $I(t)$, and preference relation, $>_t$.

An allocation $Y(t)$ is a m -tuple of vectors in Ω_d such that $\sum_{t=1}^m Y(t) \leq \sum_{t=1}^m I(t)$. A competitive equilibrium for

\mathcal{E} is a pair $\langle X, \bar{p} \rangle$ such that $\bar{p} \in \Omega_d$, $X(t)$ is an allocation with $\bar{p} \cdot X(t) \leq \bar{p} \cdot I(t)$ for all t , and such that for all $\bar{y} \in \Omega_d$ with $\bar{y} >_t X(t)$, $\bar{p} \cdot \bar{y} > \bar{p} \cdot I(t)$. $X(t)$ is called a competitive allocation and \bar{p} is called a competitive price system.

Given a representing family of utilities for a standard exchange economy \mathcal{E} , we can, of course, define a finite game in characteristic or coalitional form, where $V(S)$, the worth of a coalition, is the maximum of the sum of the individual utilities that the coalition S can obtain by redistributing their aggregate initial endowment. This game has a Shapley value which we denote as $\phi_{V, \mathcal{E}}$.

The prototypical example of an unbounded family of standard exchange economies is a sequence of replicated economies. Let \mathcal{E} be a standard exchange economy having m traders. The r -fold replica of \mathcal{E} , denoted $\mathcal{E}^{(r)}$, consists

of r copies of \mathcal{E} . That is, $\mathcal{E}^{(r)}$ with mr traders consists of m types of traders where each type has r traders and all members of a type have the same initial endowment and preference relation. Moreover, each type has the same preference and initial endowment as some trader in the original economy \mathcal{E} . Every allocation in \mathcal{E} defines an allocation in $\mathcal{E}^{(r)}$ by replication, i.e. each member of a type is given the same commodity bundle as his counterpart in \mathcal{E} . If X is an allocation in \mathcal{E} , then $X^{(r)}$ will denote the replicated allocation in $\mathcal{E}^{(r)}$.

Theorem 2. If each trader in \mathcal{E} has a smooth preference (in the sense of Debreu) and each trader has a strictly positive initial endowment, then the following are equivalent:

- (i) X is a competitive allocation for \mathcal{E} .
- (ii) $X^{(r)}$ is a competitive allocation for $\mathcal{E}^{(r)}$ for all $r \in \mathbb{N}$.
- (iii) There is a representing family of utilities $\{u_t\}_{t \in \mathcal{E}}$ with the property that for every $\epsilon > 0$ there is an $r_0 \in \mathbb{N}$ such that if $r \geq r_0$ and $S \subset \mathcal{E}^{(r)}$, then
 - (1) $|\sum_{t \in S} \varphi_{t; \mathcal{E}^{(r)}}(t) - \frac{1}{mr} \sum_{t \in S} u_t(X^{(r)}(t))| < \epsilon$.
- (iv) For every $\epsilon > 0$ and all $r_0 \in \mathbb{N}$ there is an $r > r_0$ and a representing family of utilities such that if $S \subset \mathcal{E}^{(r)}$ then Equation (1) in (iii) holds.

We shall only give an outline of the proof. Of course, (i) \Leftrightarrow (ii) is clear.

If $r_0 \in {}^*N - N$, then for any $r \geq r_0$, r_0 satisfies property (i) of Lemma 1 (i.e., $\max_{t \in S} \|X(t)\| < r_0$) for $\mathcal{E}^{(r)}$.

Therefore, there is an $r_1 \in N$ such that Lemma 2 is valid with $\gamma = r_1$ for $\mathcal{E}^{(r)}$ whenever $r \geq r_1$. Given \bar{p} and $I(t)$, we may now, as in the proof of Part a of Theorem 1, find a box $B_\gamma(0)$ and utilities $\{u_t\}$ essentially concave on $B_\gamma(0)$ so that if $\omega \in {}^*N - N$ and $\langle X^{(\omega)}, \bar{p} \rangle$ is a nonstandard competitive equilibrium for $\mathcal{E}^{(\omega)}$, then $X^{(\omega)}$ is a value allocation with respect to the family of utilities $\{u_t\}$.

The notions of competitive equilibrium which we have defined for standard and nonstandard exchange economies differ in their definition of allocation, budget set, and maximality of a commodity bundle in the budget set. But the concept of competitive equilibrium as it is defined for standard exchange economies is also meaningful in nonstandard exchange economies. We shall call this the Q-competitive equilibrium (quasi-standard competitive equilibrium).

Now fix $\epsilon > 0$ and $r_0 \in {}^*N - N$ and replicate the economy ω times where $\omega \geq r_0$. Given an allocation X in \mathcal{E} , we consider $X^{(\omega)}$. Next we use the lemma, which is proved in [4], that every Q-competitive equilibrium in $\mathcal{E}^{(\omega)}$

is a nonstandard competitive equilibrium in $\mathcal{E}^{(\omega)}$. Thus, if X is a competitive equilibrium in \mathcal{E} , then it follows immediately that $X^{(\omega)}$ is a Q -competitive equilibrium in $\mathcal{E}^{(\omega)}$. Hence, by the lemma, $X^{(\omega)}$ is a nonstandard competitive equilibrium and by the value equivalence theorem, Theorem 1, for all coalitions $S \subset \mathcal{E}^{(\omega)}$,

$$\left| \sum_{t \in S} \varphi_t V; \mathcal{E}^{(\omega)}(t) - \frac{1}{m\omega} \sum_{t \in S} u_t(X^{(\omega)}(t)) \right| < \epsilon. \text{ It follows that}$$

there is a standard $r_0 \in \mathbb{N}$ such that if $r \geq r_0$ and $S \subset \mathcal{E}^{(r)}$ then Equation 1 holds, i.e. (ii) \implies (iii).

That (iii) \implies (iv) is clear. If X satisfies (iv), then $X^{(\omega)}$ is a value allocation for some $\omega \in {}^*N - N$, i.e. choose an infinitesimal ϵ and $r_0 \in {}^*N - N$. Therefore, there exists a $\bar{p} \in {}^*\Omega_d$ such that $\langle X^{(\omega)}, \bar{p} \rangle$ is a nonstandard competitive equilibrium with $\|\bar{p}\| = 1$ and $\bar{p} \gg \bar{0}$. Since the preferences in $\mathcal{E}^{(\omega)}$ are standard, it follows that $\langle X, {}^0\bar{p} \rangle$ is a competitive equilibrium for \mathcal{E} .

REFERENCES

- [1] Aumann, R. J. "Values of Markets with a Continuum of Traders," Econometrica, 43 (1975), 611-646.
- [2] _____ and L. Shapley. Values of Non-Atomic Games. Princeton University Press, Princeton, 1974.
- [3] Brown, D. J. and A. Robinson. "Nonstandard Exchange Economies," Econometrica, 43 (1974), 41-55.
- [4] _____. "The Cores of Large Standard Exchange Economies," Journal of Economic Theory, 9 (1974), 245-254.
- [5] Brown, D. J. "Existence of a Competitive Equilibrium in a Nonstandard Exchange Economy," Econometrica (forthcoming).
- [6] Champsaur, P. "Cooperation versus Competition," unpublished manuscript, 1971.
- [7] Debreu, G. "Smooth Preferences," Econometrica, 40 (1972), 603-616.