

Yale University

## EliScholar – A Digital Platform for Scholarly Publishing at Yale

---

Cowles Foundation Discussion Papers

Cowles Foundation

---

2-1-1979

### Value Theory without Efficiency

Pradeep Dubey

Abraham Neyman

Robert J. Weber

Follow this and additional works at: <https://elischolar.library.yale.edu/cowles-discussion-paper-series>



Part of the [Economics Commons](#)

---

#### Recommended Citation

Dubey, Pradeep; Neyman, Abraham; and Weber, Robert J., "Value Theory without Efficiency" (1979).

*Cowles Foundation Discussion Papers*. 748.

<https://elischolar.library.yale.edu/cowles-discussion-paper-series/748>

This Discussion Paper is brought to you for free and open access by the Cowles Foundation at EliScholar – A Digital Platform for Scholarly Publishing at Yale. It has been accepted for inclusion in Cowles Foundation Discussion Papers by an authorized administrator of EliScholar – A Digital Platform for Scholarly Publishing at Yale. For more information, please contact [elischolar@yale.edu](mailto:elischolar@yale.edu).

COWLES FOUNDATION FOR RESEARCH IN ECONOMICS  
AT YALE UNIVERSITY

Box 2125, Yale Station  
New Haven, Connecticut 06520

COWLES FOUNDATION DISCUSSION PAPER NO. 513

NOTE: Cowles Foundation Discussion Papers are preliminary materials circulated to stimulate discussion and critical comment. Requests for single copies of a Paper will be filled by the Cowles Foundation within the limits of the supply. References in publications to Discussion Papers (other than mere acknowledgment by a writer that he has access to such unpublished material) should be cleared with the author to protect the tentative character of these papers.

VALUE THEORY WITHOUT EFFICIENCY

by

Pradeep Dubey, Abraham Neyman and Robert James Weber

February, 1979

# VALUE THEORY WITHOUT EFFICIENCY\*

by

Pradeep Dubey,<sup>1</sup> Abraham Neyman<sup>2</sup> and Robert James Weber<sup>1</sup>

## 0. Introduction

Recently attention has been focused on generalizations and analogues of the Shapley value that do not enjoy the efficiency, or Pareto optimality, property ([7], [9]). This has stemmed from the search for value functions that describe the prospects of playing different roles in a game (instead of describing fair division, in which case efficiency is a natural requirement). The purpose of this paper is to treat the subject from an axiomatic viewpoint, i.e., to characterize the class of operators that is obtained by omitting the efficiency axiom from the axioms defining the Shapley value. We consider both finite-player and nonatomic games. In the finite case, a complete solution is given; in the nonatomic case, a complete solution is given for the important space  $pNA$ .

## 1. The Finite Case

Let  $U$  be an infinite set, the universe of players. A game on  $U$  is a set function  $v : 2^U \rightarrow R$  with  $v(\emptyset) = 0$ . We interpret the members of  $U$  as players and the members of  $2^U$  as coalitions. A set  $N \subset U$

---

\*The results presented here are the intersection of work done independently by A. Neyman on the one hand, and P. Dubey and R. J. Weber on the other. This research was supported in part by grants from the National Science Foundation and the Office of Naval Research.

<sup>1</sup>Cowles Foundation for Research in Economics, Yale University.

<sup>2</sup>Department of Mathematics, University of California at Berkeley.

is a support of  $v$  if, for each  $S \subset U$ ,  $v(S) = v(S \cap N)$ . A finite game is a game which has a finite support. We denote by  $G$  the vector space of all finite games, and by  $G^N$  the subspace of  $G$  consisting of games with support  $N$ . Let  $AG$  (respectively,  $AG^N$ ) be the subspace of  $G$  (respectively,  $G^N$ ) of additive games. (Note that for  $N$  finite,  $AG^N$  is isomorphic to  $R^N$ , the Euclidean space of dimension  $|N|$  whose axes are indexed by the elements of  $N$ . For convenience we shall often use  $R^N$  for  $AG^N$ .)

Given a permutation  $\theta$  of  $U$  (i.e., a 1-1 mapping from  $U$  onto itself) define the game  $\theta*v$  by  $(\theta*v)(S) = v(\theta S)$ . Finally define  $v$  to be monotonic if  $v(S) \geq v(T)$  whenever  $S \supset T$ .

A semivalue on  $G$  is a function  $\Psi : G \rightarrow AG$  such that:

- (1)  $\Psi$  is linear,
- (2)  $\Psi\theta* = \theta*\Psi$ , for each permutation  $\theta$  of  $U$ ,
- (3) if  $v$  is monotonic, then  $\Psi v$  is monotonic,
- (4) if  $v \in AG$ , then  $\Psi v = v$ .

These are the linearity, symmetry, monotonicity and projection axioms ([1], pp. 15-16). The projection axiom is an easy consequence of the more familiar dummy axiom, which says that if  $i$  is a dummy player in  $v$  (i.e.,  $v(S \cup i) = v(S) + v(i)$  whenever  $i \notin S$ ) then  $(\Psi v)(i) = v(i)$ . (We conventionally omit the braces when indicating one-element sets.) The quantity  $(\Psi v)(i)$ , for  $i \in U$ , is a measure (according to  $\Psi$ ) of the prospect of having role  $i$  in the game  $v$ .

Let  $\xi$  be a probability measure on  $[0,1]$ . For any  $i \in U$  and any  $v \in G$  with finite support  $N$ , define  $\Psi_\xi v \in AG$  by

$$(1.1) \quad (\Psi_\xi v)(i) = \sum_{S \subset N \setminus i} p_S^\xi [v(S \cup i) - v(S)],$$

where

$$p_s^n = \int_0^1 t^s (1-t)^{n-s-1} d\xi(t) .$$

(The symbols  $n$  and  $s$  generically denote the cardinalities of the sets  $N$  and  $S$ .) Note that the right-hand side of (1.1) is independent of the choice of  $N$ , so the definition makes sense.

We now come to our characterization of semivalues on  $G$ .

Theorem 1a. For each probability measure  $\xi$  on  $[0,1]$ ,  $\Psi_\xi$  is a semivalue. Moreover, every semivalue on  $G$  is of this form, and the mapping  $\xi \rightarrow \Psi_\xi$  is 1-1.

To prove this theorem we first characterize the semivalues on the vector space of games on a fixed finite-player set. This characterization has appeared elsewhere (see, for example, [9]). For the sake of completeness, we present an alternative derivation here. Then we proceed with two different proofs which shed light on Theorem 1a from different viewpoints. Let  $N \subset U$  be a finite set. A semivalue on  $G^N$  is a function  $\psi^N : G^N \rightarrow AG^N$  satisfying (1), (2<sup>N</sup>), (3), (4), where (2<sup>N</sup>) requires that  $\psi_{\theta^*}^N = \theta^* \psi^N$  for every  $N$ -preserving permutation  $\theta$  of  $U$ .

Let  $p^n = (p_0^n, \dots, p_{n-1}^n)$  be a vector such that  $\sum_{s=0}^{n-1} \binom{n-1}{s} p_s^n = 1$  and  $p_s^n \geq 0$ . Define  $\psi_{p^n}^N : G^N \rightarrow AG$  by

$$(1.2) \quad (\psi_{p^n}^N v)(i) = \sum_{S \subset N \setminus i} p_S^n [v(S \cup i) - v(S)]$$

for all  $i \in N$  and  $v \in G^N$ .

Lemma. For each vector  $p^n$ ,  $\psi_{p^n}^N$  is a semivalue on  $G^N$ . Moreover, every semivalue on  $G^N$  is of this form, and the mapping  $p^n \rightarrow \psi_{p^n}^N$  is 1-1.

Proof. It is straightforward to verify that each  $\psi_{p^n}^N$  is indeed a semivalue. Without loss of generality take  $N = \{1, \dots, n\}$ , and let  $\psi^N$  be a semivalue on  $G^N$ . Consider the vector space  $F$  of symmetric linear functions from  $C^N$  and  $AG^N$ . For any nonempty  $S \subset N$ , define the game  $v_S \in G^N$  by  $v_S(T) = 1$  if  $S \subset T$ ,  $v_S(T) = 0$  otherwise. It is well-known (see, for example, Appendix A of [1]) that  $\{v_S : \emptyset \neq S \subset N\}$  is a basis for  $G^N$ ; therefore, every element  $f \in F$  is uniquely determined by its values on the games in this basis. From the symmetry axiom (2), it is in fact sufficient to specify  $f(v)$  for every  $v \in \{v_{S(k)} : 1 \leq k \leq n\}$  where  $S(k) = \{1, \dots, k\}$ . Hence the dimension of  $F$  is at most  $n$ .

For each  $0 \leq k \leq n-1$  let  $\Psi_{(k)} = \psi_{p^n}^N$ , as defined by (1.2) when

$p_k = \binom{n-1}{k}^{-1}$  and  $p_\ell = 0$  for all  $\ell \neq k$ . It is clear that each

$\Psi_{(k)} \in F$  and  $\{\Psi_{(0)}, \dots, \Psi_{(n-1)}\}$  is linearly independent in  $F$ . Thus this set is a basis for  $F$ .

Consider  $\psi^N \in F$ . It can be uniquely written as

$\psi^N = c_0 \Psi_{(0)} + \dots + c_{n-1} \Psi_{(n-1)}$ . Therefore we must only show that

$\sum_{s=0}^{n-1} c_s = 1$  and  $c = (c_0, \dots, c_{n-1}) \geq 0$ ; the desired result will then

follow upon taking  $p_s^n = \binom{n-1}{s}^{-1} c_s$ , yielding  $\psi^N = \psi_{p^n}^N$ . Suppose some

$c_k < 0$ . Consider  $w \in G^N$  defined by  $w(T) = 1$  if  $|T| > k$ ,  $w(T) = 0$

otherwise. Then for any  $i \in N$ ,  $(\psi^N w)(i) = c_k (\Psi_{(k)} w)(i) = c_k < 0$ ;

this contradicts the monotonicity axiom (3). Next consider  $v_{\{1\}} \in G^N$ .

By the projection axiom (4), we must have  $(\Psi^N v_{\{1\}})(1) = \sum_{s=0}^{n-1} c_s = v_{\{1\}}(1) = 1$ .  $\square$

Proof of Theorem 1a. It is straightforward to verify that each  $\Psi_\xi$  is a semivalue. Consider any semivalue  $\Psi$ . For each finite  $N \subset U$ ,  $\Psi$  induces a semivalue  $\Psi^N$  on  $G^N$ . From the preceding lemma we know that each  $\Psi^N$  has the form

$$(\Psi^N v)(i) = \bigcup_{S \subset N \setminus i} p_S^N [v(S \cup i) - v(S)]$$

where all  $p_S^N \geq 0$  and  $\sum_{s=0}^{n-1} \binom{n-1}{s} p_s^N = 1$ . Furthermore, it is a simple consequence of the symmetry axiom that there is a collection of constants  $\{p_s^n : s = 0, \dots, n-1; n = 1, 2, \dots\}$  such that for all  $i \in N \subset U$  and  $S \subset N \setminus i$ ,  $p_S^N = p_S^n$ .

Consider the collection of games  $\{\hat{v}_S^N\}$ , where  $\hat{v}_S^N$  in  $G^N$  is defined for any  $S \subset N \subset U$  by  $\hat{v}_S^N(T) = 1$  if  $T \supseteq S$ , and 0 otherwise. For any  $i \in N \setminus S$ ,

$$\Psi^N(\hat{v}_S^N)(i) = p_S^N = p_S^n.$$

For any given player  $d \in U \setminus N$ , the game  $\hat{v}_S^N$  can be viewed as a game in  $G^{N \cup d}$ . It is easily shown that for any  $i \in N \setminus S$ ,

$$\Psi^{N \cup d}(\hat{v}_S^N)(i) = p_S^{N \cup d} + p_S^{N \cup d} = p_S^{n+1} + p_{S \cup d}^{n+1}$$

Since  $\Psi^N$  and  $\Psi^{N \cup d}$  are restrictions of the same operator  $\Psi$ , it follows that for any  $i \in N \setminus S$ ,

$$(1.3) \quad \Psi^N(\hat{v}_S^N)(i) = p_S^n = p_S^{n+1} + p_{S \cup d}^{n+1} = \Psi^{N \cup d}(\hat{v}_S^N)(i).$$

For notational ease, set  $\alpha_n = p_n^{n+1}$  (for  $n = 0, 1, 2, \dots$ ). Obviously,  $p_s^n$  determines  $\{\alpha_n\}_{n=0}^{\infty}$ . Moreover, using (1.3) it can be shown by induction that for any  $0 \leq s \leq n$ ,

$$\begin{aligned} p_s^{n+1} &= (-1)^{n-s} \left( \alpha_n - \binom{n-s}{1} \alpha_{n-1} + \binom{n-s}{2} \alpha_{n-2} + \dots + (-1)^{n-s} \alpha_s \right) \\ &= (-1)^{n-s} \Delta^{n-s} \alpha_n, \end{aligned}$$

where  $\Delta$  is the standard "backwards difference" operator. Consequently, we see that every sequence  $\{\alpha_n\}$  of real numbers uniquely defines a collection  $\{p_s^n\}$ . It can be shown by direct summation that, for each  $n$ ,

the numbers  $\left\{ \binom{n-1}{s} p_s^n \right\}_{s=0}^{n-1}$  add to  $\alpha_0$ . Therefore, the collection  $\{p_s^n\}$

will define a semivalue if and only if  $\alpha_0 = 1$  and all  $p_s^n \geq 0$ .

It is well-known (for example, Theorem 4.6 of [3]) that a sequence  $\{\alpha_n\}$  (with  $\alpha_0 = 1$ ) and the successive differences  $(-1)^k \Delta^k \alpha_n$  of all orders are nonnegative if and only if  $\alpha_0, \alpha_1, \dots$  are the moments of a uniquely-determined probability distribution  $\xi$  on  $[0,1]$ . In this case, since each  $\alpha_n = \int_0^1 t^n d\xi(t)$ , it follows that each

$$\begin{aligned} p_s^{n+1} &= \int_0^1 \left( t^s - \binom{n-s}{1} t^{s+1} + \dots + (-1)^{n-s} t^n \right) d\xi(t) \\ &= \int_0^1 t^s (1-t)^{n-s} d\xi(t). \quad \square \end{aligned}$$



Alternative Proof of Theorem 1a.

It suffices to establish that  $\Psi$  is of the form  $\Psi_\xi$  for a unique probability measure  $\xi$  on  $[0,1]$ . Let  $i \in U$  be fixed. For each finite subset  $N$  of  $U \setminus i$ ,  $\Psi$  induces a semivalue on  $G^{N \cup i}$ , and hence, by Lemma 1, induces a probability measure  $c_N$  on the subsets of  $N$  such that  $c_N(S) = p_S^{n+1}$ . If  $N \subset \bar{N}$ , then by considering the natural embedding of  $G^N$  into  $G^{\bar{N}}$ , we have  $c_N(S) = \sum_{\bar{N}} c_{\bar{N}}(T)$ , where the summation runs over all  $T$  for which  $S \subset T \subset \bar{N}$  and  $T \cap N = S$ . Let  $\{N_k\}$  be an increasing sequence of finite subsets of  $U \setminus i$ . The measures on the subsets of the various  $N_k$  are "consistent," and therefore by Kolmogorov's consistency theorem ([5], p. 94), there is a sequence of  $(0,1)$ -valued random variables  $\{Y_j : j \in \cup N_k\}$  such that  $c_{N_k}(S) = \text{Prob}(\{j : Y_j = 1\} = S)$ . Thus  $\{Y_j\}$  is an exchangeable sequence of random variables. De Finetti's theorem ([4], sec. 9.6.1) asserts that the distribution of every exchangeable infinite sequence of random variables is a unique mixture of distributions of sequences of independent identically-distributed random variables. As  $\text{Prob}(Y_j = 0 \text{ or } 1) = 1$ , there exists a unique probability measure  $\xi$  on  $[0,1]$  such that for every finite sequence  $\{\epsilon_j : j \in N\}$  of 0's and 1's,  $\text{Prob}(Y_j = \epsilon_j \text{ for all } j \in N) = \int_0^1 t^{\sum \epsilon_j} (1-t)^{n-\sum \epsilon_j} d\xi(t) = c_N(\{j : \epsilon_j = 1\})$ .

It is obvious from the axiom of symmetry that the mixing measure

depends neither on the particular player  $i$ , nor on the sequence  $N_k$ , and thus  $\xi$  is uniquely determined by  $\Psi$  alone.  $\square$

This alternative proof provides another view of the theorem. Let  $(\Omega, \mathcal{B}, P)$  be a probability space, and  $\{X_i : i \in U\}$  a family of independent identically-distributed random variable distributed uniformly on  $[0,1]$ . If  $v \in G$  and  $t \in [0,1]$ , define the random variable  $\Delta v(t)$  by  $\Delta v(t) = v(\{i : X_i \leq t\}) - v(\{i : X_i < t\})$ . We then have the following restatement of Theorem 1a:

Theorem 1a'. For each probability measure  $\xi$  on  $[0,1]$  there is a semi-value  $\Psi_\xi$  on  $G$  defined by

$$(\Psi_\xi v)(i) = \int_0^1 E(\Delta v(t) | X_i = t) \cdot d\xi(t) .$$

Moreover, every semivalue on  $G$  is of this form and the mapping  $\xi \rightarrow \Psi_\xi$  is 1-1 .

The Shapley value [8] is defined as  $\phi = \Psi_\lambda$ , where  $\lambda$  denotes the Lebesgue measure on  $[0,1]$ . This is the only semivalue which has the efficiency property: for every  $N \subset U$  and  $v \in G^N$ ,  $\phi v(N) = v(N)$ . Define the bounded-variation norm of a game  $v \in G$  with support  $N$ , as  $\|v\| = \inf(v_+(N) + v_-(N))$ , where the infimum is taken over all pairs  $v_+, v_-$  of monotonic games for which  $v = v_+ - v_-$ . With respect to this norm on  $G$ , the Shapley value is a continuous linear operator of norm 1. (For any monotonic  $v_+, v_- \in G^N$  such that  $v = v_+ - v_-$ ,  $\|\phi v\| = \sum |\phi v(i)| \leq \sum (\phi v_+(i) + \phi v_-(i)) = v_+(N) + v_-(N)$ ; hence  $\|\phi v\| \leq \|v\|$ . But for any monotonic  $v \in G^N$ ,  $\|\phi v\| = v(N) = \|v\|$ .)

We shall characterize the class of continuous semivalues on  $G$ .

Let  $W$  be the subset of  $L_\infty(0,1)$  of all nonnegative functions  $g$  with  $\int_0^1 g(t)dt = 1$ .

Theorem 1b. For each  $g \in W$ , the operator  $\Psi_g : G \rightarrow AG$  defined by

$$\Psi_g v(i) = \int_0^1 E(\Delta v(t) | X_i = t) \cdot g(t) dt$$

is a continuous semivalue. Moreover, every continuous semivalue on  $G$  is of this form. The map  $g \rightarrow \Psi_g$  is a linear isometry (that is,  $\|\Psi_g\| = \|g\|_{L_\infty}$ ).

Proof. Consider any  $g \in W$  and define  $\xi = \int g d\lambda$ . By Theorem 1a',  $\Psi_g = \Psi_\xi$  is a semivalue. For any  $v \in G^N$ , and monotonic games  $v_+, v_-$  with  $v = v_+ - v_-$ ,  $\|\Psi_g v\| = \sum |\Psi_g v(i)| \leq \sum |\Psi_g v_+(i)| + \sum |\Psi_g v_-(i)| \leq \|g\| \cdot (\sum |\phi v_+(i)| + \sum |\phi v_-(i)|) = \|g\| \cdot (v_+(N) + v_-(N))$ ; therefore,  $\|\Psi_g v\| \leq \|g\| \cdot \|v\|$ . Hence  $\Psi_g$  is continuous, and  $\|\Psi_g\| \leq \|g\|$ .

Next, consider any continuous semivalue  $\Psi_\xi$ . Select any (relatively) open interval  $J \subset [0,1]$ , and assume that  $\xi(J) = M \cdot \lambda(J)$ . Fix a player  $i \in U$ , and for each  $k > 0$ , select  $N_k \subset U$  such that  $i \in N_k$  and  $|N_k| = k$ . Let  $v_k \in G^{N_k}$  be defined by  $v_k(S) = \lambda([0, \frac{S}{n}] \cap J)$ . By the law of large numbers,  $\liminf \Psi_\xi v_k(i) \geq \frac{1}{n} \cdot \xi(J) = \frac{M}{n} \cdot \lambda(J)$ . Therefore  $\|\Psi_\xi v_k\| = \sum |\Psi_\xi v_k(i)| \geq M \cdot \lambda(J)$ , while each  $\|v_k\| = \lambda(J)$ . Hence,  $\|\Psi_\xi\| \geq M$ . The continuity of  $\Psi_\xi$  implies that  $\|\Psi_\xi\|$  is finite. Consequently,  $\bar{M} = \sup\{\xi(J)/\lambda(J) : J \text{ is an interval in } [0,1]\} < \infty$ , and the Radon-Nikodym derivative  $d\xi/d\lambda = g$  is in  $W$ . Therefore  $\Psi_\xi = \Psi_g$ , and  $\|\Psi_g\| \geq \bar{M} = \|g\|$ .  $\square$

## 2. The Infinite Case

All definitions and notation are according to [1]. Let  $(I, C)$  be a measure space isomorphic to  $([0,1], B)$ , where  $B$  is the  $\sigma$ -field of Borel subsets of  $[0,1]$ . The members of  $I$  are called players, the members of  $C$  coalitions, and set functions are called games. Let  $BV$  be the space of bounded-variation set functions on  $(I, C)$ . The space of all bounded, finitely-additive set functions is denoted  $FA$ , and its subspace of all nonatomic measures is denoted  $NA$ . Denote by  $G$  the group of automorphisms of  $(I, C)$ . For each  $\theta \in G$ ,  $\theta^* : BV \rightarrow BV$  is defined by  $\theta^*v(S) = v(\theta S)$ . If  $Q \subset BV$  then  $Q^+$  denotes the subset of  $Q$  of all monotonic set functions. A subset  $Q$  of  $BV$  is symmetric if for each  $\theta \in G$ ,  $\theta^*Q \subset Q$ . An operator  $\psi : Q \rightarrow BV$  is called positive if  $\psi(Q^+) \subset BV^+$ , and symmetric if for each  $\theta \in G$ ,  $\theta^*\psi = \psi\theta^*$ .

Let  $Q$  be a linear symmetric subspace of  $BV$ . A semivalue on  $Q$  is an operator  $\psi$  from  $Q$  into  $FA$  such that:

- (1)  $\psi$  is linear,
- (2)  $\psi$  is symmetric,
- (3)  $\psi$  is positive,
- (4) if  $v \in Q \cap FA$  then  $\psi v = v$ .

We will characterize the semivalues on  $pNA$ , the closed subspace of  $BV$  spanned by all powers of  $NA^+$  measures. This space plays an important role in the theory of nonatomic games, and contains many games of interest. For example,  $pNA$  contains all "vector measure games" satisfying appropriate differentiability conditions, i.e., all set functions of the form  $f \circ \mu$ , where  $\mu = (\mu_1, \dots, \mu_n)$  is a nonatomic finite-dimensional vector measure and  $f$  is an appropriately differentiable real-valued function defined on the range of  $\mu$ , with  $f(0) = 0$ . As

our main theorem in this section uses notation and terminology related to the "extension" of a game, we restate here relevant definitions and results from [1].  $I$  denotes the family of all measurable functions from  $(I, \mathcal{C})$  to  $([0,1], \mathcal{B})$ . There is a partial order on  $I$ :  $f \geq g$  if  $f(s) \geq g(s)$  for all  $s \in I$ . A real valued function  $w$  on  $I$  with  $w(0) = 0$  is called an ideal set function; it is called monotonic if  $f \geq g$  implies  $w(f) \geq w(g)$ . The characteristic function of a member  $S$  of  $\mathcal{C}$  is denoted  $\chi_S$ . We will sometimes denote  $\chi_S$  by  $S$  and  $t \cdot \chi_I$  by  $t$ .

It is shown in [1; Theorem G] that there is a unique monotonicity-preserving linear mapping which associates with each  $v \in \text{pNA}$  an ideal set function  $v^*$ , such that  $(v \cdot w)^* = v^* \cdot w^*$  for all  $v, w \in \text{pNA}$ , and  $\mu^*(f) = \int_I f \cdot d\mu$  for all  $\mu \in \text{NA}$  and  $f \in I$ .

Denote  $\partial v^*(t, S) = (d/d\tau) \cdot v^*(t\chi_I + \tau\chi_S) \Big|_{\tau=0}$ . By Theorem H of [1] we know that for each  $v \in \text{pNA}$  and each  $S \in \mathcal{C}$ , the derivative  $\partial v^*(t, S)$  exists for almost all  $t$  in  $[0,1]$ , and is integrable over  $[0,1]$  as a function of  $t$ .

Recall that  $W$  is the set of nonnegative functions  $g \in L_\infty(0,1)$  such that  $\int_0^1 g(t) dt = 1$ .

Theorem 2. For each  $g \in W$  the operator  $\Psi_g : \text{pNA} \rightarrow \text{FA}$  defined by

$$\Psi_g v(S) = \int_0^1 \partial v^*(t, S) \cdot g(t) dt$$

is a semivalue. Moreover, every semivalue on  $\text{pNA}$  is of this form. The map  $g \rightarrow \Psi_g$  of  $W$  onto the family of semivalues on  $\text{pNA}$  is a linear isometry.

Proof. Let  $g \in W$  be given. For  $v \in \text{pNA}$ , Lemma 23.1 of [1] asserts

that  $\int_0^1 |\partial v^*(t, S)| \cdot dt \leq \|v\|$ . Hence  $|\Psi_g v(S)| = \left| \int_0^1 \partial v^*(t, S) \cdot g(t) dt \right| \leq \|g\| \cdot \|v\|$ ;

this proves that  $\Psi_g v$  is bounded. If  $S, T \subset I$  with  $S \cap T = \emptyset$  then

$\partial v^*(t, T \cup S) = \partial v^*(t, T) + \partial v^*(t, S)$  for almost all  $t$ . Therefore

$\Psi_g v(S \cup T) = \Psi_g v(S) + \Psi_g v(T)$ , which proves that  $\Psi_g$  takes  $\text{pNA}$  into  $\text{FA}$ .

Linearity of  $\Psi_g$  follows from the linearity of the extension as well as

that of the derivative. Symmetry of  $\Psi_g$  follows from the fact that

$\partial(\theta^*v)^*(t, S) = \partial v^*(t, \theta S)$  and thus  $\theta^* \Psi_g v(S) = \int \partial v^*(t, \theta S) \cdot g(t) dt$

$= \int \partial(\theta^*v)^*(t, S) \cdot g(t) dt = \Psi_g(\theta^*v)(S)$ . Let  $v \in \text{pNA}^+$ . Then  $v^*$  is also

monotonic and  $\partial v^*(t, S) \geq 0$ ; thus  $\Psi_g v$  is monotonic, which proves the

positivity of  $\Psi_g$ . Finally, any  $u \in \text{pNA} \cap \text{FA}$  is in  $\text{NA}$  (Corollary

5.3 of [1], and the continuity of the elements of the space  $\text{AC}$  ([1],

page 205), imply that  $u$  is countably additive). Hence  $\partial u^*(t, S) = u(S)$

and consequently  $\Psi_g u = u$ . This completes the proof that  $\Psi_g$  is a semi-

value.

Now, let  $\Psi$  be a semivalue on  $\text{pNA}$ . Let  $\mu$  be a fixed probability measure in  $\text{NA}$ . Each  $f \in L_1$  induces a game  $v_f$  defined by

$$v_f(S) = \int_0^{\mu(S)} f(t) dt.$$

In other words,  $f$  defines a function  $F: [0, 1] \rightarrow \mathbb{R}$  by  $F(s) = \int_0^s f(t) dt$ ,

and  $v_f = F \circ \mu$ . As  $f \in L_1$ ,  $F$  is absolutely continuous and therefore

$v_f \in \text{pNA}$ . In analogy with the proof of Proposition 6.1 of [1] it follows

that  $\Psi v_f = C(f) \cdot \mu$ , where  $C(f)$  is a constant independent of  $\mu$ . Ob-

serve that  $v_{f+g} = v_f + v_g$ ; thus the linearity of  $\Psi$  implies that  $C$

is linear. We now proceed to show that  $C$  is continuous. Observe that

$\|v_f\| = \|f\|_{L_1}$ . Since  $pNA$  is internal ([1], Proposition 7.19), it is a closed reproducing space and thus ([1], Proposition 4.3)  $\Psi$  is continuous on  $pNA$ . That is, there exists a constant  $K$  with  $\|\Psi v\| \leq K \cdot \|v\|$ , which in particular implies that  $\|C(f)\| = \|C(f) \cdot \mu\| \leq K \cdot \|v_f\| = K \cdot \|f\|_{L_1}$ .

Hence  $C : L_1 \rightarrow \mathbb{R}$  is a continuous linear functional and therefore is of the form  $C(f) = \int_0^1 f(t) g(t) dt$  for some  $g \in L_\infty$ . We shall show that

$\Psi = \Psi_g$ . As was shown in the beginning of the proof,  $\Psi_g(pNA) \subset FA$  and  $|\Psi_g v(S)| \leq \|g\| \cdot \|v\|$ , which implies that  $\Psi_g$  is continuous. For each  $f \in L_1$ ,  $\partial v_f^*(t, S) = f(t) \cdot \mu(S)$  for almost all  $t$ , and thus

$$\Psi_g v_f(S) = \mu(S) \cdot \int f(t) g(t) dt = C(f) \cdot \mu(S) = \Psi v_f(S) \quad \text{and therefore} \quad \Psi_g v_f = \Psi v_f.$$

The linear symmetric subspace spanned by  $\{v_f : f \in L_1\}$  is dense in  $pNA$  (it contains all powers of  $NA$  measures). The operators  $\Psi$  and  $\Psi_g$  are linear and symmetric and thus coincide on this subspace; as they are also continuous, they coincide on  $pNA$ . It remains for us to show that

$g \in W$ . For  $v \in NA \subset FA \cap pNA$ , it follows that  $\partial v^*(t, S) = v(S)$ .

Thus  $\Psi_g v(S) = \left(\int_0^1 g(t) dt\right) v(S)$ , which shows that  $\int_0^1 g(t) dt = 1$ . Let

$B_\epsilon = \{t : g(t) \leq -\epsilon\}$  and let  $f$  be the characteristic function of  $B_\epsilon$ .

Then  $f \geq 0$  and hence  $v_f$  is monotonic. But as  $\Psi_g v_f(I) = \int f(t) g(t) dt \leq -\epsilon \cdot \lambda(B_\epsilon)$  ( $\lambda$  denotes the Lebesgue measure on  $[0, 1]$ ) and  $\Psi_g = \Psi$

is positive, it must be that  $\lambda(B_\epsilon) = 0$ . As this holds for any  $\epsilon > 0$ ,

$g$  is nonnegative. This completes the proof that any semivalue  $\Psi$  is of the form  $\Psi_g$  for some  $g \in W$ .

Now, for any  $g \in W$  and  $\epsilon > 0$  there exists a nonnegative  $f \in L_1$  with

$\|f\|_{L_1} = 1$  and  $\int f(t) g(t) dt = \|g\| - \epsilon$ . Observe that  $\|v_f\| = \|f\|_{L_1} = 1$  and that

$\|\Psi_g v_f\| = \|g\| - \epsilon$ ; hence  $\|\Psi_g\| \geq \|g\|$ . On the other hand, for  $v \in pNA^+$ ,

$$\|\Psi_g v\| = \Psi_g v(I) = \int \partial v^*(t, I) \cdot g(t) dt \leq \|g\| \cdot \int_0^1 \partial v^*(t, I) \cdot dt = \|g\| \cdot \|v\| .$$

In the general case, when  $v$  is not necessarily monotonic, let  $\epsilon > 0$  be given. Set  $v = u - w$ , where  $u$  and  $w$  are in  $pNA^+$  and

$$\|v\| + \epsilon \geq \|u\| + \|w\| ; \text{ such } u \text{ and } w \text{ exist because } pNA \text{ is internal.}$$

$$\text{Then } \|\Psi_g v\| \leq \|\Psi_g u\| + \|\Psi_g w\| \leq \|g\|(\|u\| + \|w\|) \leq \|g\|(\|v\| + \epsilon) ,$$

and if we let  $\epsilon \rightarrow 0$ ,  $\|\Psi_g v\| \leq \|g\| \cdot \|v\|$ ; this completes the proof of the equality  $\|\Psi_g\| = \|g\|$ .  $\square$

### 3. Remarks

Continuous semivalues are diagonal. (The proof in [6] that continuous values are diagonal does not make use of the efficiency axiom and therefore the same proof works here.) Furthermore, semivalues on closed reproducing spaces are diagonal.

The semivalues derived axiomatically on  $pNA$  can also be obtained from a complementary, asymptotic point of view [2] which links the finite-player and nonatomic approaches.



## REFERENCES

- [1] R. J. Aumann and L. S. Shapley, Values of Nonatomic Games, Princeton University Press, 1974.
- [2] P. Dubey, Asymptotic semivalues and a short proof of Kannai's theorem, Math. of Oper. Res., to appear.
- [3] S. Karlin and W. J. Studden, Tchebycheff Systems, with Applications in Analysis and Statistics, John Wiley & Sons, 1966.
- [4] B. de Finetti, Probability, Induction and Statistics, John Wiley & Sons, 1972.
- [5] M. Loève, Probability Theory, D. van Nostrand Co., 1955.
- [6] A. Neyman, Continuous values are diagonal, Math. of Oper. Res. 2 (1977), 338-342.
- [7] A. E. Roth, The Shapley value as a von Neumann-Morgenstern utility, Econometrica 45 (1977), 657-664.
- [8] L. S. Shapley, A value for n-person games, Ann. Math. Study 28 (1953), 307-318.
- [9] R. J. Weber, Probabilistic values for games, Cowles Foundation Discussion Paper No. 471R, Yale University, 1978.