

Yale University

EliScholar – A Digital Platform for Scholarly Publishing at Yale

Public Health Theses

School of Public Health

January 2022

Residential And Race/ethnicity Disparities In Heat Vulnerability In The United States

Mitchell Manware
memanware@gmail.com

Follow this and additional works at: <https://elischolar.library.yale.edu/ysphtdl>

Recommended Citation

Manware, Mitchell, "Residential And Race/ethnicity Disparities In Heat Vulnerability In The United States" (2022). *Public Health Theses*. 2304.
<https://elischolar.library.yale.edu/ysphtdl/2304>

This Open Access Thesis is brought to you for free and open access by the School of Public Health at EliScholar – A Digital Platform for Scholarly Publishing at Yale. It has been accepted for inclusion in Public Health Theses by an authorized administrator of EliScholar – A Digital Platform for Scholarly Publishing at Yale. For more information, please contact elischolar@yale.edu.

Residential and Race/Ethnicity Disparities in Heat Vulnerability in the United States

Name: Mitchell Manware

Year Completed: 2022

Year Degree Awarded: 2023

Degree Awarded: Master of Public Health

Department: School of Public Health

Advisor: Dr. Kai Chen

Committee Member(s): Dr. Yusuf Ransome

Abstract

Adverse health outcomes caused by extreme heat represent the most direct human health threat associated with the warming of the Earth's climate. Socioeconomic, demographic, health, land cover, and temperature determinants contribute to heat vulnerability; however, nationwide patterns of residential and race/ethnicity disparities in heat vulnerability in the United States are poorly understood. This study aimed to develop a Heat Vulnerability Index (HVI) for the United States; to assess differences in heat vulnerability across geographies that have experienced historical and/or contemporary forms of marginalization; and to quantify HVI by race/ethnicity. Principal components analysis (PCA) was used to calculate census tract level HVI scores based on the 2019 population characteristics of the United States. Differences in HVI scores were analyzed across the Home Owners' Loan Corporation (HOLC) "redlining" grades, the Climate and Economic Justice Screening Tool (CEJST) disadvantaged versus non-disadvantaged communities, and race/ethnicity groups. HVI scores were calculated for 55,267 U.S. census tracts. Mean HVI scores were 17.56, 18.61, 19.45, and 19.93 for HOLC grades 'A' – 'D,' respectively. CEJST-defined disadvantaged census tracts had a significantly higher mean HVI score (19.13) than non-disadvantaged tracts (16.68). The non-Hispanic African American or Black race/ethnicity group had the highest HVI score (18.51), followed by Hispanic or Latino (18.19). Historically redlined and contemporary CEJST disadvantaged census tracts and communities of color were found to be associated with increased vulnerability to heat. These findings can help promote equitable climate change adaptation policies by informing policymakers about the national distribution of place- and race/ethnicity-based disparities in heat vulnerability.

Acknowledgements

We gratefully acknowledge support from the Yale Planetary Solutions Project seed grant and the High Tide Foundation.

Authors

Mitchell Manware, mitchell.manware@yale.edu.^{1,2}

Dr. Robert Dubrow, robert.dubrow@yale.edu.^{2,3}

Dr. Daniel Carrión, daniel.carrion@yale.edu.^{2,3}

Yiqun Ma, yiqun.ma@yale.edu.^{2,3}

Dr. Kai Chen, kai.chen@yale.edu.^{2,3}

¹Department of Social and Behavioral Sciences, Yale School of Public Health, New Haven, Connecticut, USA

²Yale Center on Climate Change and Health, Yale School of Public Health, New Haven, Connecticut, USA.

³Department of Environmental Health Sciences, Yale School of Public Health, New Haven, Connecticut, USA.

Conflict of Interest

The authors declare no conflicts of interest relevant to this study.

Data Availability Statement

Data on the heat vulnerability index and its indicators will be publicly available after the publication at https://github.com/CHENlab-Yale/HVI_US.

Table of Contents

1. Introduction
2. Methods
 - 2.1. Data sources
 - 2.1.1. Heat Vulnerability Index (HVI) Data
 - 2.1.2. Residential Disparity Data
 - 2.2. Statistical Analysis
 - 2.2.1. HVI Construction
 - 2.2.2. Heat Vulnerability and Environmental Racism Analysis
3. Results
4. Discussion
5. Conclusion
6. Tables
 - 6.1. Table 1 Heat Vulnerability Index Variables and Supporting Literature
 - 6.2. Table 2 Heat vulnerability index data sources and descriptive statistics
7. Figures
 - 7.1. Figure 1 Heat Vulnerability Index Score Maps
 - 7.2. Figure 2 Distribution of Census Tract Heat Vulnerability Index Scores for HOLC Grades
 - 7.3. Figure 3 Heat Vulnerability Index Scores for Disadvantaged Communities and HOLC Grades
 - 7.4. Figure 4 Heat Vulnerability Index Scores for Race/Ethnicity Groups
8. References

1. Introduction

Adverse health outcomes associated with extreme heat represent the most direct human health threat of the warming of the Earth's climate caused by anthropogenic greenhouse gas emissions (Xu et al., 2016; Zhai et al., 2021). Disparities in heat exposure, exacerbated by disparities in access to individual- and community-level protective factors, result in inequitable distribution of the burden of heat-related illness across geographic locations, occupations, and demographic characteristics (Campbell et al., 2018). The disproportionate exposure of communities of color in the United States to environmental hazards, often ascribed at least in part to racially-driven residential segregation policies, is a fundamental manifestation of environmental racism (Bravo et al., 2016; Chavis, 1994; Gee & Payne-Sturges, 2004; Lopez, 2002; Morello-Frosch & Jesdale, 2006; Perry, 2007). Recently, heat-related outpatient emergency department visits and inpatient admissions from the emergency department were found to be higher in “redlined” neighborhoods, evidence of the potential adverse health consequences at the intersection of racist policies and climate change (Li et al., 2021).

The term “redlining” stems from lending maps created by the Home Owners’ Loan Corporation (HOLC) in the 1930s (Percy, 2020). Across urban areas in the United States, maps were created to assess a neighborhood’s level of loan risk based on its physical building conditions, public utilities and, infamously, demographic composition (Percy, 2020). Neighborhoods with African American, Hispanic, Asian, and Jewish populations were often graded the lowest, resulting in denied mortgage loans and limited wealth accumulation among these communities (Aaronson et al., 2021; Percy, 2020). City-wide maps displaying neighborhood grades were color coded, with the highest-risk areas colored red, giving rise to the term “redlining” (Percy, 2020). Recently, the White House Council on Environmental Quality created the Climate and Economic Justice Screening Tool (CEJST), which identifies “disadvantaged” communities based on various socioeconomic, environmental, and health criteria (*Climate and Economic Justice Screening Tool*). The CEJST provides a contemporary measure of residential disparity in the United States, though differing from HOLC grades by its inclusion of suburban and rural areas, as well as by its equity promoting purpose (*Climate and Economic Justice Screening Tool*).

HOLC “redlining” grades and CEJST advantage status provide unique, insightful measures of residential disparity, each supporting environmental justice research by capturing a range of place-based exposures and outcomes (Smoyer, 1998). Previous studies have observed higher land surface temperature, less green space, and more dark roofs in “redlined” census tracts (Hoffman et al., 2020; 2021; Namin et al., 2020; Nardone et al., 2021; Schinasi et al., 2022; Wilson, 2020). However, these studies did not assess the association between HOLC grade and various socioeconomic, demographic, or biological determinants of heat vulnerability (Organization, 2021), nor did they analyze potential differences in heat vulnerability by race/ethnicity. Furthermore, previous heat vulnerability indices have been developed for individual cities (Conlon et al., 2020; Johnson et al., 2012; Mallen et al., 2019; Wolf & McGregor, 2013), counties (Harlan et al., 2013; Prudent et al., 2016), or states (Maier et al., 2014; Nayak et al., 2018), and the only national heat vulnerability index was limited by missing data and lack of a heat exposure variable (Reid et al., 2009).

To fill these gaps, we aimed to better understand place-based and race/ethnicity-based disparities in heat vulnerability across the entire United States. We build on the aforementioned research by 1) developing an expansive and uniform Heat Vulnerability Index (HVI) for the contiguous United States; 2) assessing how the HVI differs across historical (i.e., HOLC grade) and contemporary (i.e., CEJST) measures of residential disparity; and 3) quantifying the HVI distribution by race/ethnicity. We hypothesized that heat vulnerability would be greatest in both historically-defined and contemporaneously-defined disadvantaged communities and in communities of color.

2. Methods

2.1 Data Sources

2.1.1 Heat Vulnerability Index (HVI) Data

We evaluated the peer-reviewed literature to identify candidate variables for the HVI, (i.e., variables found to be associated with heat-related adverse health outcomes; **Table 1**). Of the variables with supporting evidence, only those with nationwide data available at the county or census tract levels were included. Data used to construct the HVI included demographic, economic, social, housing, diabetes, land cover, and temperature variables collected from various

sources (**Table 2**). All data points were collected at or calculated to the census tract level. We obtained complete data for 55,267 census tracts, with 18,735 census tracts excluded from the HVI analysis due to missing data for one or more variables.

Demographic, economic, social and housing data was obtained at the census tract level from various American Community Survey (ACS) 5-Year Estimates from 2019. Demographic data consisted of the Hispanic or Latino, non-Hispanic African American or Black, and elderly population proportions of each census tract (*ACS Demographic and Housing Estimates*, 2019). Economic data included the proportion of unemployed individuals and individuals living below the federal poverty level (*Poverty Status in the Past 12 Months*, 2019; *Selected Economic Characteristics*, 2019). Social variables included the proportions of individuals living alone, elderly and living alone, disabled, with less than a high school education, born in a foreign country, and who are Limited English Proficient (LEP) (*Selected Social Characteristics in the United States*, 2019). The proportion of homes built prior to 1980 constituted the housing data (*Physical Housing Characteristics for Occupied Housing Units*, 2019).

Diabetes prevalence was used to represent underlying health conditions based on the association between diabetes and heat-related adverse health outcomes (Knowlton et al., 2009; Schwartz, 2005; Semenza et al., 1999; Semenza et al., 1996), the co-prevalence of diabetes with several other comorbid conditions (Iglay et al., 2016), and the availability of nationwide diabetes prevalence data (Zou et al., 2017). Sex-specific census-tract-level diabetes prevalence for persons aged 20 years and older was estimated using county-level diabetes data collected from the Center for Disease Control and Prevention (CDC) U.S. Diabetes Surveillance System and census-tract-level age and sex data collected from the ACS 5-Year Age and Sex Estimates (*Age and Sex*, 2019; Zou et al., 2017). Specifically, we used CDC sex-specific diabetes prevalence estimates for individuals aged 20 years and older for years 2015 to 2019 to calculate a 5-year, sex-specific estimate for each county. Then, we multiplied each sex- and county-specific prevalence estimate by the number of males and females aged 20 years and older, respectively, in each census tract within each county to estimate the total number of diabetic male and female individuals in each census tract. The sex-specific totals were summed and divided by the total

population aged 20 years and older in each census tract to estimate the total 5-year average diabetes prevalence in each census tract.

We obtained two land cover variables -- high density development and non-green space -- from the National Neighborhood Data Archive (NaNDA): Land Cover by Census Tract, United States, 2001-2016 data file (Clarke & Melendez, 2019). High density development is an NaNDA-defined variable that measures the proportion of land within each census tract having urban imperviousness greater than 79% (Clarke & Melendez, 2019). The non-green space variable was calculated by summing the proportion of land within each census tract that was covered in “green space” (deciduous forest, evergreen forest, mixed forest, shrub/scrub, herbaceous, hay/pasture, cultivated crops, woody wetlands, emergent herbaceous wetlands), subtracting from 1 and multiplying by 100 (Clarke & Melendez, 2019). This value represents the percentage of land in each census tract covered by non-natural, artificial surfaces (Clarke & Melendez, 2019).

A third land cover variable -- an average Enhanced Vegetation Index (EVI) score, which is an alternative measure of vegetation greenness -- was calculated for each census tract using data downloaded from NASA’s MOD13A2 Version 6 Product (Didan, 2015). A four-year average of summer-month EVI scores was calculated using 16-day EVI scores from June 1 through August 31 for years 2017 through 2020 (Didan, 2015). One km² grid cell layers representing each 16-day period within the date range were stacked to calculate the average EVI score for each grid cell (Didan, 2015). The average EVI values for each grid cell were aggregated to the census tract level by taking the area-weighted mean of each EVI score within the census tract polygon (Didan, 2015). This calculation yielded a census-tract-level average EVI score for the summer months from 2017 to 2020 (Didan, 2015). The Enhanced Vegetation Index was preferred over the Normalized Difference Vegetation Index (NDVI) for its increased sensitivity over heavily vegetated areas and correction for both atmospheric conditions and canopy background (Nawrot et al., 2007; Qian et al., 2008).

Air temperature at two-meter height was obtained from NASA’s Daymet Version 4 dataset and used to measure heat exposure for each census tract (Thornton et al., 2020). A four-year

average of non-missing air temperatures was calculated for each one km² grid cell using air temperature data from June 1 through August 31 for years 2017 through 2020 (Thornton et al., 2020). Average grid cell air temperature was aggregated to the census tract level by taking the area-weighted mean of each air temperature value within the census tract polygon, yielding average two-meter air temperature for the summer months. Census tracts' most frequently occurring temperature (MFT) was calculated using air temperature at two-meter height data from January 1, 2000 through December 31, 2020, obtained from NASA's Daymet Version 4 dataset (Thornton et al., 2020). Temperature estimates were rounded to the nearest degree, and the modal temperature estimate for each grid cell represented the MFT. Grid cell MFT estimates were aggregated to the census tract level by taking the area-weighted mean of each temperature estimate within the census tract polygon, yielding the 20-year MFT for each census tract. Heat exposure for each census tract was measured by calculating the difference between the average 2-meter air temperature for the summer months and the 20-year MFT. Quantifying heat exposure based on recent summer months' temperature anomaly relative to the MFT captures spatial differences in summertime warming and potentially incorporates an aspect of local climate adaptation (Spangler & Wellenius, 2020).

2.1.2 Residential Disparity Data

Home Owners Loan Corporation (HOLC) data were obtained from the Inter-university Consortium for Political and Social Research's *Historic Redlining Scores for 2010 and 2020 US Census Tracts* project (Meier & Mitchell, 2021). This project overlaid HOLC maps with 2020 census tract polygons to calculate a weighted average HOLC score for recent census tract boundaries (Meier & Mitchell, 2021). HOLC grades ranged from "A" to "D," where 'A' was "Best," 'B' was "Desirable," 'C' was "Definitely Declining," and 'D' was "Hazardous" (Nelson et al., 2020; Percy, 2020). The original HOLC grades were converted to numbers for calculation purposes (A=1, B=2, C=3, D=4) (Meier & Mitchell, 2021). The resulting census-tract-level-weighted HOLC scores for 2020 census tract boundaries ranged continuously from 1.0 to 4.0 and were available for 13,488 census tracts across 142 cities (Meier & Mitchell, 2021).

Disadvantaged community data were obtained from the White House Council on Environmental Quality's Climate and Economic Justice Screening Tool (CEJST) (*Climate and*

Economic Justice Screening Tool). Data utilized for this analysis consisted of the tool’s binary census-tract-specific disadvantage indicator (“disadvantaged” or “non-disadvantaged”), along with each census tract’s unique geographic identifier (GEOID) (*Climate and Economic Justice Screening Tool*). The CEJST identifies a census tract as “disadvantaged” if it exceeds the threshold for one or more environmental or climate indicators and for both socioeconomic indicators within at least one of the tool’s eight categories of criteria (*Climate and Economic Justice Screening Tool*).

Race and ethnicity population data were collected from American Community Survey 5-Year Demographic and Housing Estimates from 2019 (*ACS Demographic and Housing Estimates*, 2019). Race and ethnicity were combined, producing eight groups for comparison: Hispanic or Latino of any race, non-Hispanic White, non-Hispanic Black or African American, non-Hispanic American Indian and Alaska Native, non-Hispanic Asian, non-Hispanic Native Hawaiian and Other Pacific Islander, non-Hispanic Other race, and non-Hispanic Multi-racial (two or more races).

2.2 Statistical Analysis

2.2.1 HVI Construction

Principal component analysis (PCA) was performed to reduce the original number of variables to a smaller number of sufficiently explanatory components. PCA reduces potential collinearity between original variables by grouping correlated variables into new principal components (Reid et al., 2009). The PCA method has been used previously to construct city- (Conlon et al., 2020; Johnson et al., 2012; Mallen et al., 2019; Wolf & McGregor, 2013), county- (Harlan et al., 2013; Prudent et al., 2016), state- (Maier et al., 2014; Nayak et al., 2018), and national- (Reid et al., 2009) level HVIs. The varimax rotation method minimizes the number of variables loading on specific principal components, ensuring they are independent of each other and the original variables (Reid et al., 2009). Principal components are retained if they meet each of the following standard criteria: a clear break in the Scree plot, Eigenvalue greater than 1, and the cumulative proportion of variance explained by the retained components is greater than 70% (Cattell, 1966; Jolliffe & Cadima, 2016; Wold et al., 1987). Census tracts receive normalized

factor scores for each retained component, which are often rescaled for easier interpretation and to minimize the effect of outliers in subsequent calculations (Reid et al., 2009).

We calculated an HVI for 55,267 census tracts, with 18,735 census tracts excluded from the analysis due to missing data for one or more variables. Together, the EVI and air temperature datasets contributed 15,927 (90.7%) of the missing data points, possibly due to grid cell disruption caused by water body or cloud coverage. Correlation among the included variables was assessed using Spearman's rank correlation coefficient (**Table S1** in the Supplemental Information). PCA with varimax rotation identified five components to be retained based on meeting the standard criterion, adequately representing the various aspects of heat vulnerability (**Figures S1-S2** in the Supplemental Information). Factor loadings quantify each variable's influence on the individual retained components, and can be described as Education/Language, Economic/Demographic, Land Cover/Housing, Elderly/Social Isolation, and Exposure/Underlying Conditions factors (**Table S2** in the Supplemental Information). For interpretability and calculation purposes, normalized factor scores produced from the varimax rotation were rescaled from 1 to 6 based on deviation from the mean. Each census tract, therefore, received five factor scores, each ranging from 1 to 6. The individual factor scores were summed to calculate the HVI score for each census tract, having a potential range of 5 to 30. HVI scores were geocoded and plotted using each census tract's unique GEOID.

2.2.2 Heat Vulnerability and Environmental Racism Analysis

Correlation and linear regression analysis methods were used to quantify the relationship between HVI scores and weighted HOLC scores at the census tract level. Weighted HOLC scores were rounded to the nearest integer and reassigned original grade measurements (1=A, 2=B, 3=C, 4=D) to assess differences in HVI score among grade-specific strata. We used analysis of variance to test whether at least one HOLC grade-specific mean HVI score was significantly different from at least one other grade-specific score. If this was the case, we then used post-hoc honestly significant difference tests to calculate the magnitude, direction, and statistical significance of differences between HOLC grade-specific mean HVI scores. We compared mean HVI scores for CEJST "disadvantaged" and "non-disadvantaged" census tracts using the t-test. We used analysis of variance and post-hoc honestly significant difference tests to

assess differences in mean HVI score among HOLC-grade specific census tract strata by CEJST disadvantaged community status.

Heat vulnerability was analyzed for the various race/ethnicity groups in the United States in several ways. First, a mean HVI score was calculated for each group weighted by the census-tract-specific population of each group. In addition, each census tract was assigned to a race/ethnicity category according to its modal, or most populous, race or ethnicity. Mean census tract HVI scores were then compared across race/ethnicity categories. For example, a census tract where 2,000 residents were Hispanic or Latino of any race, 1,000 were non-Hispanic White, 500 were non-Hispanic African American or Black, and 500 were non-Hispanic Asian, would be assigned to the Hispanic or Latino of any race category. The mean HVI score of all census tracts assigned to the Hispanic or Latino of any race category would be calculated and compared to that of other race/ethnicity categories. Finally, the population proportion of each race/ethnicity group was calculated for HVI score deciles.

3. Results

HVI scores were calculated for 55,267 U.S. census tracts (**Figure 1**). Scores ranged from 10 to 26 with a country-wide mean (SD) equal to 17.47 (2.16). Visual assessment of the HVI Map revealed higher HVI scores in urban census tracts compared to surrounding suburban and rural areas. In the greater New York City metropolitan area, higher HVI scores were apparent in the Bronx, Brooklyn and Newark, New Jersey compared to lower scores in the western portion of Nassau County, New York, and Bergen County in northeastern New Jersey. In Chicago, high HVI scores were evident throughout Cook County, with lower scores more prevalent in the surrounding DuPage and Will Counties. Los Angeles and Houston's urban areas also exhibited considerably higher HVI scores than the surrounding suburbs. **Figure 1** also depicts higher HVI scores in the southwestern United States, a spatial pattern consistent with the Global Burden of Disease (GBD) estimates of heat-related mortality at the state level (**Figure S3** in the Supplemental Information) (*Global Burden of Disease Study 2019 (GBD 2019) Results*, 2020). A sensitivity analysis was conducted to compare the original HVI scores, which used the average summertime 2m air temperature anomaly relative to the full-year MFT as the temperature anomaly variable, to alternative HVI scores using the average summertime 2m air temperature anomaly

relative to the summertime (June-August) MFT as the temperature anomaly variable. The sensitivity analysis yielded similar HVI results, with the original and alternative HVI scores highly correlated ($R = 0.88$) (**Table S3 and Figure S4** in the Supplemental Information).

U.S. Department of Agriculture's Rural-Urban Commuting Area codes were used to assess differences in mean HVI scores between urban and non-urban areas (*2010 Rural-Urban Commuting Area Codes*, 2019). T test analysis measured significantly different (p value < 0.001) mean HVI score in urban areas compared to non-urban areas. On average, HVI scores in urban areas were 0.29 (0.25, 0.33) points higher than in non-urban areas, supporting the visual differences apparent in **Figure 1**.

There was a weak-to-moderate association between HOLC grade and HVI (correlation coefficient of 0.305). Based on a simple linear model, per unit increase in weighted HOLC score was associated with a 0.8-point increase in HVI score (**Figure S5** in the Supplemental Information). Mean (SD) HVI scores were 17.56 (1.80) for HOLC grade A census tracts, 18.61 (2.08) for grade B, 19.45 (1.92) for grade C, and 19.93 (1.90) for grade D (**Figure 2**). These mean HVI scores differed significantly across HOLC grade strata (p value < 0.001). Post-hoc honestly significant difference tests calculated significantly higher (p value < 0.001) HVI scores for each incrementally higher-risk HOLC grade, differing by 1.05 (0.81, 1.28) between 'A' and 'B,' by 0.84 (0.71, 0.97) between 'B' and 'C,' and by 0.48 (0.37, 0.60) between 'C' and 'D' (**Figure 2**).

We assessed the distribution of the raw HVI variables across the various HOLC grades using heat maps (**Figure S6** in the Supplemental Information). Analysis of variance showed that the mean value of each raw HVI variable significantly differed across HOLC grades (**Table S4** in the Supplemental Information), suggesting that each raw variable contributed to the difference in mean HVI scores across the HOLC grade strata.

The mean (SD) HVI score for CEJST-defined "disadvantaged" census tracts was 19.13 (1.89), significantly higher ($p < 0.001$) than the mean HVI score of 16.68 (1.80) for "non-disadvantaged" census tracts. We also calculated the mean HVI score for census tracts within

each HOLC grade stratified by disadvantage community status and found the highest score for disadvantaged census tracts with HOLC grade ‘D’ and the lowest for non-disadvantaged tracts with HOLC grade ‘A’ (**Figure 3**). Analysis of variance showed that mean HVI scores significantly differed across HOLC grades among both the “disadvantaged” ($p < 0.001$) and “non-disadvantaged” ($p < 0.001$) census tracts. Incrementally higher-risk HOLC grades had significantly higher ($p < 0.001$) HVI score among non-disadvantaged census tracts. However, the only significant difference among disadvantaged tracts was between grade ‘C’ and grade ‘D.’ Within each HOLC grade stratum, disadvantaged census tracts had a significantly higher ($p < 0.001$) mean HVI score than non-disadvantaged tracts.

The mean (SD) HVI score was highest for the non-Hispanic Black or African American race/ethnicity group at 18.51 (2.02), followed by 18.19 (2.29) for Hispanic or Latino of any race, 17.95 (2.29) for Other races, 17.31 (2.44) for Asian, 17.05 (1.90) for American Indian or Alaskan Native, 17.05 (2.14) for Multi-race, 16.76 (1.90) for non-Hispanic White, and 16.76 (2.13) for Native Hawaiian or Pacific Islander. These mean HVI scores differed significantly ($p = 0.036$) across groups. When calculated based on the modal race/ethnicity group population in each census tract, the mean (SD) HVI score was highest for Other races at 22.0 (0.0) (based on only two census tracts in which Other races had the modal population), followed by 19.31 (1.70) for non-Hispanic Black or African American, 19.06 (2.14) for Hispanic or Latino of any race, 18.51 (2.66) for Asian, 16.96 (1.71) for American Indian or Alaskan Native, and 16.90 (1.89) for non-Hispanic White (**Figure 4a**). The Native Hawaiian or Pacific Islander and Multi-race race/ethnicity groups were not the modal population in any census tract.

Non-white persons were overrepresented in the most heat-vulnerable census tracts (**Figure 4b**). Non-white individuals made up 75.6% of the population in the highest HVI score category of census tracts, compared to only 24.7% in the lowest category. Hispanic or Latino and non-Hispanic African American or Black individuals accounted for 38.4% and 26.6% of the highest category population, respectively, compared to 10.7% and 4.3% in the lowest category population and 18.0% and 12.3% in the entire United States population (*ACS Demographic and Housing Estimates*, 2019).

4. Discussion

The analyses in this study each supported the hypothesis that vulnerability to heat-related adverse health outcomes is greatest in historically-defined and contemporaneously-defined disadvantaged communities and in communities of color in the United States. Historically “redlined” census tracts were associated with higher heat vulnerability, consistent with similar analyses that found higher land surface temperature and lower green space in these segregated areas (Hoffman et al., 2020; Li et al., 2021; Namin et al., 2020; Nardone et al., 2021; Schinasi et al., 2022; Wilson, 2020). This association was evidenced by the correlation and regression analyses between continuous, weighted HOLC score and HVI score, as well as significantly increasing HVI scores between incrementally higher-risk HOLC grades. These residential disparities were driven by inequitably distributed determinants of heat vulnerability (i.e., the raw HVI variables), each of which differed significantly among HOLC grade strata.

Contemporaneously-defined disadvantaged community status was also found to be associated with higher HVI score, both in aggregate and across HOLC grade strata. When HOLC grade and CEJST disadvantaged community status were assessed together, the highest place-based heat vulnerability was measured in census tracts that were both historically redlined and presently disadvantaged. Finally, we observed high mean HVI scores among non-White race/ethnicity groups, and overrepresentation of people of color in the most vulnerable census tracts. Together, these results demonstrate place- and race/ethnicity-based disparities in heat vulnerability across the United States, and add to the literature on the long-term adverse environmental and health implications associated with racist housing policies (Hoffman et al., 2020; Li et al., 2021; Namin et al., 2020; Nardone et al., 2021; Schinasi et al., 2022; Wilson, 2020).

In addition to residential and race/ethnicity differences in heat vulnerability, urban areas were measured to have higher HVI scores than non-urban areas. This difference in vulnerability, however, cannot be validated without nationwide estimates of heat-related mortality. Although previous heat vulnerability indices also have measured increased vulnerability in urban areas (Reid et al., 2009), other studies have identified higher risk of heat mortality in rural areas (Chen et al., 2016; Hu et al., 2019). Heat vulnerability in urban versus non-urban communities has

complex differences; therefore adaptation strategies should be designed to target the specific needs of each area.

The HVI developed in this study utilizes methodologies applied in previous studies (Reid et al., 2009), but also builds on previous indices in several ways. First, constructing the index with data from the most recently updated datasets provides more temporally relevant measures of the included variables. Utilizing the most recent temperature estimates in particular is crucial for adequately describing heat exposure, as warming trends differ between the most and least socially vulnerable counties in the United States (Spangler & Wellenius, 2020). Second, the air temperature variable used to construct our HVI measures heat exposure more accurately (Organization, 2021) than does land surface temperature (LST), which has been used in the construction of previous indices. LST likely misrepresents heat exposure, as atmospheric and surface conditions can cause several degree differences between LST and ambient air temperature (Good, 2016). Additionally, measuring heat exposure based on recent summer months' temperature anomaly relative to the MFT captures current and spatially diverse warming trends. Basing the temperature anomaly on MFT, which is closely associated with minimum mortality temperature, also helps take into account potential geography-specific temperature adaptation, making it a multifaceted measure of heat exposure (Yin et al., 2019). Finally, this assessment provides HVI scores for most United States' census tracts ($n = 55,267$; 75% of census tracts), capturing vulnerability across urban, suburban, and rural areas. An earlier study sought to measure heat vulnerability for the entire country but excluded many non-urban areas from the final vulnerability scores due to the lack of a nationwide dataset for air conditioning (Reid et al., 2009). Other studies of heat vulnerability within individual cities (Conlon et al., 2020; Johnson et al., 2012; Mallen et al., 2019; Wolf & McGregor, 2013), counties (Harlan et al., 2013; Prudent et al., 2016), or states (Maier et al., 2014; Nayak et al., 2018) did not provide the national scope necessary for an environmental racism assessment.

Beyond the improvements to the HVI, this study provided a unique analysis of the place- and race/ethnicity-based disparities associated with heat vulnerability. Previous studies have measured higher land surface temperature and lower green space in “redlined” areas (Hoffman et al., 2020; Li et al., 2021; Namin et al., 2020; Nardone et al., 2021; Schinasi et al., 2022; Wilson,

2020), but these studies failed to capture the various other socioeconomic, demographic or biological determinants of vulnerability (Organization, 2021). Additionally, this study is the first to quantify heat vulnerability across the various race/ethnicity groups in the United States, further illustrating the vulnerability of communities of color.

This study had several limitations. First, 18,735 census tracts did not receive an HVI score due to missing data and were therefore excluded from the analysis. Data loss occurred primarily during the calculation of census tract level EVI and air temperature values, possibly due to grid cell disruption caused by cloud coverage (Shen et al., 2015). We cannot rule out the possibility of some selection bias due to exclusion of these census tracts. A second limitation was the lack of a nationwide dataset measuring census-tract-level air conditioning access, which is known to provide strong protection against heat-related morbidity and mortality (Bouchama et al., 2007; Kaiser et al., 2001; Kilbourne et al., 1982; Naughton et al., 2002; Petkova et al., 2014; Romanello et al., 2021; Semenza et al., 1996). The positive association that has been observed between warmer temperatures and air conditioning access could indicate more widespread access to air conditioning across the southern portion of the country, further influencing the spatial distribution of HVI scores across the United States if air conditioning access had been included as a variable in the HVI construction (Bell et al., 2009). A third limitation was the use of diabetes to represent all comorbidities and underlying health conditions associated with heat-related morbidity and mortality. In fact, the magnitude of the association between heat exposure and various physical and mental health conditions varies (Knowlton et al., 2009; Semenza et al., 1999; Semenza et al., 1996); however, measures of underlying health conditions other than diabetes were unavailable nationally at the county- or census-tract-level. Additionally, the original variables used to construct the HVI were consistent across geographic locations, but heat vulnerability in urban versus non-urban areas could be dependent on which combination of raw variables are used (Fard et al., 2021). Though a potential limitation, the observed urban versus non-urban differences in heat vulnerability are likely driven by differences in exposure, susceptibility, and adaptability that are captured in the original variables.

A fourth limitation of this study was that we were unable to validate the HVI by comparison to actual empirical heat-related morbidity or mortality data, due to data unavailability, but the

general spatial pattern of HVI scores had approximate consistency with the GBD estimates of heat-related mortality at the state level (*Global Burden of Disease Study 2019 (GBD 2019) Results*, 2020). A further limitation was that the Home Owners' Loan Corporation assigned grades based on the demographic, economic and social construction of a neighborhood, implicitly taking into account many of the same variables used for construction of the HVI score (Nardone et al., 2021; Percy, 2020). Finally, the White House Council on Environmental Quality's Climate and Economic Justice Screening Tool, which was utilized for this analysis due to its national scope, did not include race/ethnicity as a criterion for identifying environmentally and socially disadvantaged communities in the United States. Race/ethnicity also is not taken into account in California's CalEnviroScreen (*CalEnviroScreen 4.0*, 2021), but is a factor in the New York State Disadvantage Communities Map (*Disadvantaged Communities Map*). Due to varying geographic domains, a unique combination of demographic, socioeconomic and environmental determinants, as well as different development methodologies, comparing our HVI scores to disadvantaged community status defined by these other tools would result in different outcomes and conclusions.

5. Conclusion

Residential and racial inequalities associated with heat vulnerability identified by this study confirm the persistent legacy of racist policies in the United States and their connection to contemporary environmental injustices. Our findings can help inform policymakers about the national distribution of place- and race-based disparities in heat vulnerability to develop equity-promoting climate adaptation policies.

6. Tables

Table 1 Heat Vulnerability Index Variables and Supporting Literature^a

Variable	Supporting Literature
Percentage population that is Hispanic or Latino	Uejio et al. (2011); Knowlton et al. (2009); Jones et al. (1982); Lin et al. (2009)
Percentage population that is non-Hispanic African American or Black	Uejio et al. (2011); Schwartz (2005); Whitman et al. (1997); Jones et al. (1982); O'Neill et al. (2005); O'Neill et al. (2003); Zanobetti et al. (2013)
Percentage population that is elderly	Gronlund et al. (2016); Semenza et al. (1999); Anderson and Bell (2009); Jones et al. (1982); Whitman et al. (1997)
Percentage population that is unemployed	Anderson and Bell (2009); Hansen et al. (2013)
Percentage population living below the federal poverty level	Curriero et al. (2002); Naughton et al. (2002); Jones et al. (1982); Zanobetti et al. (2013)
Percentage population that is living alone	Naughton et al. (2002); Semenza et al. (1996); Uejio et al. (2011)
Percentage population that is elderly and living alone	Gronlund et al. (2016); Semenza et al. (1999); Anderson and Bell (2009); Jones et al. (1982); Whitman et al. (1997); Naughton et al. (2002); Semenza et al. (1996); Uejio et al. (2011)
Percentage population that is disabled	Bark (1998); Bark (1982); Kaiser et al. (2001)
Percentage population with less than a high school education	Curriero et al. (2002); Zanobetti et al. (2013); O'Neill et al. (2003)
Percentage population that was born in a foreign country	Chow et al. (2012)
Percentage population that is Limited English Proficient (LEP)	Uejio et al. (2011); McGeehin and Mirabelli (2001); Hansen et al. (2013)
Percentage of occupied houses built before 1980	Gronlund et al. (2015); Xu et al. (2013)
Percentage of adults age 20+ diagnosed with diabetes	Semenza et al. (1999); Semenza et al. (1996); Schwartz (2005); Knowlton et al. (2009)
Percentage of land covered in high density development	Uejio et al. (2011); Jones et al. (1982); Harlan et al. (2006); Clarke (1972); (Buechley et al., 1972)
Percentage of land covered in non-green space	Gronlund et al. (2015); Zanobetti et al. (2013); Kilbourne et al. (1982); Xu et al. (2013); Harlan et al. (2006); Harlan et al. (2013)
Average summertime Enhanced Vegetation Index (EVI) score	Gronlund et al. (2015); Harlan et al. (2006); Harlan et al. (2013)
Average summertime 2m air temperature anomaly (°F)	Harlan et al. (2006); Johnson and Wilson (2009); Avashia et al. (2021); Basu et al. (2008); Medina-Ramón and Schwartz (2007); Zanobetti and Schwartz (2008)

^a Supporting literature identifies research studies that support a relationship between the given variables and adverse heat-related health outcomes.

Table 2 Heat vulnerability index data sources and descriptive statistics

Category	Data Source	Year Range	Variable	Mean	SD	(Min, Max)	
Demographic variables	American Community Survey 5-Year Demographic and Housing Estimates	2015 - 2019	Percentage population that is Hispanic or Latino	16.58	21.44	(0, 100)	
		2015 - 2019	Percentage population that is non-Hispanic African American or Black	14.08	22.25	(0, 100)	
		2015 - 2019	Percentage population that is elderly	16.54	7.91	(0, 92.2)	
	American Community Survey 5-Year Selected Economic Characteristics	2015 - 2019	Percentage population that is unemployed	3.54	2.49	(0, 100)	
		2015 - 2019	Percentage population that is living alone	28.23	11.26	(0, 100)	
		2015 - 2019	Percentage population that is elderly and living alone	11.29	6.11	(0, 100)	
		American Community Survey 5-Year Selected Social Characteristics	2015 - 2019	Percentage population that was born in a foreign country	12.58	13.59	(0, 100)
			2015 - 2019	Percentage population that is Limited English Proficient (LEP)	7.88	10.69	(0, 100)
			2015 - 2019	Percentage population that is disabled	11.18	6.33	(0, 100)
			2015 - 2019	Percentage population with less than a high school education	12.76	10.35	(0, 100)
American Community Survey 5-Year Poverty Status in the Past 12 Months	2015 - 2019	Percentage population living below the federal poverty level	14.75	11.59	(0, 100)		
Housing variable	American Community Survey 5-Year Physical Housing Characteristics for Occupied Housing Unit	2015 - 2019	Percentage of occupied houses built before 1980	60.94	26.19	(0, 100)	
Underlying health conditions variable	Centers for Disease Control and Prevention. U.S. Diabetes Surveillance System website	2014 - 2018	Percentage of adults age 20+ diagnosed with diabetes	6.83	1.38	(0.99, 15.78)	
	American Community Survey 5-Year Age and Sex	2015 - 2019					
Land cover variables	National Neighborhood Data Archive (NaNDA): Land Cover by Census Tract, United States, 2001-2016	2016	Percentage of land covered in high density development	9.2	15.51	(0, 100)	
		2016	Percentage of land covered in non-green space	64.89	36.66	(0.03, 100)	
	MOD13A2 MODIS/Terra Vegetation Indices	2017 - 2020	Average summertime Enhanced Vegetation Index (EVI) score	0.37	0.14	(0, 0.68)	

	16-Day L3 Global 1km SIN Grid V006‡					
Exposure variable	Daymet: Daily Surface Weather Data on a 1-km Grid for North America, Version 4	2017 - 2020	Average summertime 2m air temperature anomaly (°F)	2.97	6.09	(-4.51, 40.54)
‡ NASA's Terra and Aqua satellites' Moderate Resolution Imaging Spectroradiometer(<i>About</i>)						

7. Figures

Figure 1 Heat Vulnerability Index Score Maps

N = 55,267 census tracts. Census tracts with missing data for one or more variables were excluded from the analysis and did not receive an HVI score (N=18,735; white color on the map). Enhanced Vegetation Index (EVI) and air temperature were the primary contributors to missing data.




Figure 2 Distribution of Census Tract Heat Vulnerability Index Scores for HOLC Grades
“HOLC Grade” x-axis label represents Home Owner’s Loan Corporation lending risk grade. Grade D represents historically “redlined” census tracts. Differences in HVI scores between adjacent HOLC grades were statistically significant (p value < 0.001) according to post-hoc honest significant difference tests. Boxes depict HVI score interquartile range for each HOLC grade stratum. The middle line within the box marks the median HVI score for each HOLC grade. $N = 10,701$ census tracts with both a HOLC grade and complete data to calculate an HVI score





Figure 3 Heat Vulnerability Index Scores for Disadvantaged Communities and HOLC Grades



Violin plots show the density of census tracts receiving a particular HVI score given their HOLC grade and disadvantaged community status. Wider portions of the violin indicate more census tracts with the corresponding HVI score, while narrower portions indicate fewer census tracts with corresponding HVI score. The black dot and error bar within each violin mark the mean HVI score and standard deviation for each HOLC grade and disadvantaged community status stratum. N = 10,701 census tracts with both a HOLC grade and complete data to calculate an HVI score.



Mean (SD) Heat Vulnerability Index Scores					
	A	B	C	D	ANOVA p-value
<i>Disadvantaged Community</i>	19.8 (1.3)	20.1 (1.8)	20.2 (1.7)	20.4 (1.8)	< 0.001
<i>Non-Disadvantaged Community</i>	17.3 (1.7)	17.8 (1.8)	18.4 (1.7)	18.8 (1.7)	< 0.001

Figure 4 Heat Vulnerability Index Scores for Race/Ethnicity Groups

a) Boxes depict HVI score interquartile range for each race/ethnicity group. The middle line within the box marks the median HVI score. The Native Hawaiian or Pacific Islander and Multi-race groups were not included because neither was the modal group in any census tract. b) Bar plot depicts the race/ethnicity group population proportions within each HVI score category. Category “1” represents the lowest HVI scores and category “7” represents the highest HVI scores. These are the HVI score ranges corresponding to each category: category 1: 10 – 15 (N = 9,497; 18.6%); category 2: 16 (N = 8,313; 16.3%); category 3: 17 (N = 9,380; 18.4%); category 4: 18 (N = 8,935; 17.5%); category 5: 19 (N = 6,600; 12.9%); category 6: 20 (N = 4,180; 8.2%); category 7: 21 – 26 (N = 4,124; 8.1%). N = census tracts.



8. References

- 2010 Rural-Urban Commuting Area Codes. (2019). [CSV Data File].
<https://www.ers.usda.gov/data-products/rural-urban-commuting-area-codes/>
- Aaronson, D., Hartley, D., & Mazumder, B. (2021). The effects of the 1930s HOLC “redlining” maps. *Am Econ J Econ Policy*, 13(4), 355-392. <https://doi.org/10.1257/pol.20190414>
- About. National Aeronautics and Space Administration. <https://modis.gsfc.nasa.gov/about/>
- ACS Demographic and Housing Estimates. (2019). [CSV Data File]. U.S. Census Bureau.
https://data.census.gov/cedsci/table?q=demographic&g=0100000US%241400000&tid=A_CSDP5Y2019.DP05
- Age and Sex. (2019). [CSV Data File]. U.S. Census Bureau.
<https://data.census.gov/cedsci/table?q=age%20and%20sex&g=0100000US%241400000&tid=ACSST5Y2019.S0101>
- Anderson, B. G., & Bell, M. L. (2009). Weather-related mortality: how heat, cold, and heat waves affect mortality in the United States. *Epidemiology*, 20(2), 205-213.
<https://doi.org/10.1097/EDE.0b013e318190ee08>
- Avashia, V., Garg, A., & Dholakia, H. (2021). Understanding temperature related health risk in context of urban land use changes. *Landscape Urban Plann*, 212, 104107.
<https://doi.org/10.1016/j.landurbplan.2021.104107>
- Bark, N. (1998). Deaths of psychiatric patients during heat waves. *Psychiatr Serv*, 49(8), 1088-1090. <https://doi.org/10.1176/ps.49.8.1088>
- Bark, N. M. (1982). The prevention and treatment of heatstroke in psychiatric patients. *Hosp Community Psychiatry*, 33(6), 474-476. <https://doi.org/10.1176/ps.33.6.474>
- Basu, R., Feng, W. Y., & Ostro, B. D. (2008). Characterizing temperature and mortality in nine California counties. *Epidemiology*, 19(1), 138-145.
<https://doi.org/10.1097/EDE.0b013e31815c1da7>
- Bell, M. L., Ebisu, K., Peng, R. D., & Dominici, F. (2009). Adverse health effects of particulate air pollution: modification by air conditioning. *Epidemiology*, 20(5), 682-686.
<https://doi.org/10.1097/EDE.0b013e3181aba749>
- Bouchama, A., Dehbi, M., Mohamed, G., Matthies, F., Shoukri, M., & Menne, B. (2007). Prognostic factors in heat wave-related deaths: a meta-analysis. *Arch Intern Med*, 167(20), 2170-2176. <https://doi.org/10.1001/archinte.167.20.ira70009>
- Bravo, M. A., Anthopoulos, R., Bell, M. L., & Miranda, M. L. (2016). Racial isolation and exposure to airborne particulate matter and ozone in understudied US populations: environmental justice applications of downscaled numerical model output. *Environ Int*, 92-93, 247-255. <https://doi.org/10.1016/j.envint.2016.04.008>
- Buechley, R. W., Van Bruggen, J., & Truppi, L. E. (1972). Heat island equals death island? *Environ Res*, 5(1), 85-92. [https://doi.org/10.1016/0013-9351\(72\)90022-9](https://doi.org/10.1016/0013-9351(72)90022-9)
- CalEnviroScreen 4.0. (2021, Oct 20, 2021). Office of Environmental Health Hazard Assessment. <https://oehha.ca.gov/calenviroscreen/report/calenviroscreen-40>
- Campbell, S., Remenyi, T. A., White, C. J., & Johnston, F. H. (2018). Heatwave and health impact research: a global review. *Health Place*, 53, 210-218.
<https://doi.org/10.1016/j.healthplace.2018.08.017>
- Cattell, R. B. (1966). The scree test for the number of factors. *Multivariate Behav Res*, 1(2), 245-276. https://doi.org/10.1207/s15327906mbr0102_10
- Chavis, B. F. (1994). Preface. In R. D. Bullard (Ed.), *Unequal Protection : Environmental Justice and Communities of Color* (pp. xi - xii). Sierra Club Books.

- Chen, K., Zhou, L., Chen, X., Ma, Z., Liu, Y., Huang, L., Bi, J., & Kinney, P. L. (2016). Urbanization level and vulnerability to heat-related mortality in Jiangsu Province, China. *Environ Health Perspect*, 124(12), 1863-1869. <https://doi.org/10.1289/EHP204>
- Chow, W. T. L., Chuang, W.-C., & Gober, P. (2012). Vulnerability to extreme heat in metropolitan Phoenix: spatial, temporal, and demographic dimensions. *Professional Geographer*, 64(2), 286-302. <https://doi.org/10.1080/00330124.2011.600225>
- Clarke, J. F. (1972). Some effects of the urban structure on heat mortality. *Environ Res*, 5(1), 93-104. [https://doi.org/10.1016/0013-9351\(72\)90023-0](https://doi.org/10.1016/0013-9351(72)90023-0)
- Clarke, P., & Melendez, R. (2019). *National Neighborhood Data Archive (NaNDA): Land Cover by Census Tract, United States, 2001 - 2016 Version 1*) Inter-university Consortium for Political and Social Research. <https://doi.org/10.3886/E110663V1>
- Climate and Economic Justice Screening Tool*. Council on Environmental Quality. <https://screeningtool.geoplatform.gov/en/methodology#3/33.47/-97.5>
- Climate and Economic Justice Screening Tool*. [CSV Data File]. <https://screeningtool.geoplatform.gov/en/downloads>
- Conlon, K. C., Mallen, E., Gronlund, C. J., Berrocal, V. J., Larsen, L., & O'Neill, M. S. (2020). Mapping human vulnerability to extreme heat: a critical assessment of heat vulnerability indices created using principal components analysis. *Environ Health Perspect*, 128(9), 097001. <https://doi.org/10.1289/EHP4030>
- Curriero, F. C., Heiner, K. S., Samet, J. M., Zeger, S. L., Strug, L., & Patz, J. A. (2002). Temperature and mortality in 11 cities of the eastern United States. *Am J Epidemiol*, 155(1), 80-87. <https://doi.org/10.1093/aje/155.1.80>
- Didan, K. (2015). *MOD13A2 MODIS/Terra Vegetation Indices 16-Day L3 Global 1km SIN Grid V006* [Dataset]. NASA EOSDIS Land Processes DAAC. <https://doi.org/10.5067/MODIS/MOD13A2.006>
- Disadvantaged Communities Map*. New York State Department of Environmental Conservation. <https://climate.ny.gov/Our-Climate-Act/Disadvantaged-Communities-Criteria/Disadvantaged-Communities-Map>
- Fard, B. J., Mahmood, R., Hayes, M., Rowe, C., Abadi, A. M., Shulski, M., Medcalf, S., Lookadoo, R., & Bell, J. E. (2021). Mapping heat vulnerability index based on different urbanization levels in Nebraska, USA. *GeoHealth*, 5(10), e2021GH000478. <https://doi.org/https://doi.org/10.1029/2021GH000478>
- Gee, G. C., & Payne-Sturges, D. C. (2004). Environmental health disparities: a framework integrating psychosocial and environmental concepts. *Environ Health Perspect*, 112(17), 1645-1653. <https://doi.org/10.1289/ehp.7074>
- Global Burden of Disease Study 2019 (GBD 2019) Results*. (2020). Institute for Health Metrics and Evaluation (IHME). <https://vizhub.healthdata.org/gbd-results/>
- Good, E. J. (2016). An in situ-based analysis of the relationship between land surface “skin” and screen-level air temperatures. *J Geophys Res Atmos*, 121(15), 8801-8819. <https://doi.org/10.1002/2016JD025318>
- Gronlund, C. J., Berrocal, V. J., White-Newsome, J. L., Conlon, K. C., & O'Neill, M. S. (2015). Vulnerability to extreme heat by socio-demographic characteristics and area green space among the elderly in Michigan, 1990-2007. *Environ Res*, 136, 449-461. <https://doi.org/10.1016/j.envres.2014.08.042>

- Gronlund, C. J., Zanobetti, A., Wellenius, G. A., Schwartz, J. D., & O'Neill, M. S. (2016). Vulnerability to renal, heat and respiratory hospitalizations during extreme heat among U.S. elderly. *Clim Change*, 136(3), 631-645. <https://doi.org/10.1007/s10584-016-1638-9>
- Hansen, A., Bi, L., Saniotis, A., & Nitschke, M. (2013). Vulnerability to extreme heat and climate change: is ethnicity a factor? *Glob Health Action*, 6, 21364. <https://doi.org/10.3402/gha.v6i0.21364>
- Harlan, S. L., Brazel, A. J., Prashad, L., Stefanov, W. L., & Larsen, L. (2006). Neighborhood microclimates and vulnerability to heat stress. *Soc Sci Med*, 63(11), 2847-2863. <https://doi.org/10.1016/j.socscimed.2006.07.030>
- Harlan, S. L., Declet-Barreto, J. H., Stefanov, W. L., & Petitti, D. B. (2013). Neighborhood effects on heat deaths: social and environmental predictors of vulnerability in Maricopa County, Arizona. *Environ Health Perspect*, 121(2), 197-204. <https://doi.org/10.1289/ehp.1104625>
- Hoffman, J. S., Shandas, V., & Pendleton, N. (2020). The effects of historical housing policies on resident exposure to intra-urban heat: a study of 108 US urban areas. *Climate*, 8(1), 12. <https://doi.org/10.3390/cli8010012>
- Hu, K., Guo, Y., Hochrainer-Stigler, S., Liu, W., See, L., Yang, X., Zhong, J., Fei, F., Chen, F., Zhang, Y., Zhao, Q., Chen, G., Chen, Q., Zhang, Y., Ye, T., Ma, L., Li, S., & Qi, J. (2019). Evidence for urban-rural disparity in temperature-mortality relationships in Zhejiang Province, China. *Environ Health Perspect*, 127(3), 037001. <https://doi.org/doi:10.1289/EHP3556>
- Igley, K., Hannachi, H., Joseph Howie, P., Xu, J., Li, X., Engel, S. S., Moore, L. M., & Rajpathak, S. (2016). Prevalence and co-prevalence of comorbidities among patients with type 2 diabetes mellitus. *Curr Med Res Opin*, 32(7), 1243-1252. <https://doi.org/10.1185/03007995.2016.1168291>
- Johnson, D. P., Stanforth, A., Lulla, V., & Lubert, G. (2012). Developing an applied extreme heat vulnerability index utilizing socioeconomic and environmental data. *Appl Geogr*, 35(1), 23-31. <https://doi.org/10.1016/j.apgeog.2012.04.006>
- Johnson, D. P., & Wilson, J. S. (2009). The socio-spatial dynamics of extreme urban heat events: The case of heat-related deaths in Philadelphia. *Appl Geography*, 29(3), 419-434. <https://doi.org/10.1016/j.apgeog.2008.11.004>
- Jolliffe, I. T., & Cadima, J. (2016). Principal component analysis: a review and recent developments. *Philos Trans Royal Soc A*, 374(2065), 1-16. <https://doi.org/10.1098/rsta.2015.0202>
- Jones, T. S., Liang, A. P., Kilbourne, E. M., Griffin, M. R., Patriarca, P. A., Wassilak, S. G., Mullan, R. J., Herrick, R. F., Donnell, H. D., Jr., Choi, K., & Thacker, S. B. (1982). Morbidity and mortality associated with the July 1980 heat wave in St Louis and Kansas City, Mo. *JAMA*, 247(24), 3327-3331. <https://www.ncbi.nlm.nih.gov/pubmed/7087075>
- Kaiser, R., Rubin, C. H., Henderson, A. K., Wolfe, M. I., Kieszak, S., Parrott, C. L., & Adcock, M. (2001). Heat-related death and mental illness during the 1999 Cincinnati heat wave. *Am J Forensic Med Pathol*, 22(3), 303-307. <https://doi.org/10.1097/00000433-200109000-00022>
- Kilbourne, E. M., Choi, K., Jones, T. S., & Thacker, S. B. (1982). Risk factors for heatstroke: a case-control study. *JAMA*, 247(24), 3332-3336. <https://doi.org/10.1001/jama.1982.03320490030031>

- Knowlton, K., Rotkin-Ellman, M., King, G., Margolis, H. G., Smith, D., Solomon, G., Trent, R., & English, P. (2009). The 2006 California heat wave: impacts on hospitalizations and emergency department visits. *Environ Health Perspect*, *117*(1), 61-67. <https://doi.org/10.1289/ehp.11594>
- Li, D., Newman, G. D., Wilson, B., Zhang, Y., & Brown, R. D. (2021). Modeling the relationships between historical redlining, urban heat, and heat-related emergency department visits: An examination of 11 Texas cities. *Environ Plann Urban Anal City Sci*, *49*(3), 933-952. <https://doi.org/10.1177/23998083211039854>
- Lin, S., Luo, M., Walker, R. J., Liu, X., Hwang, S. A., & Chinery, R. (2009). Extreme high temperatures and hospital admissions for respiratory and cardiovascular diseases. *Epidemiology*, *20*(5), 738-746. <https://doi.org/10.1097/EDE.0b013e3181ad5522>
- Lopez, R. (2002). Segregation and black/white differences in exposure to air toxics in 1990. *Environ Health Perspect*, *110*(Suppl 2), 289-295. <https://doi.org/10.1289/ehp.02110s2289>
- Maier, G., Grundstein, A., Jang, W., Li, C., Naeher, L. P., & Shepherd, M. (2014). Assessing the performance of a vulnerability index during oppressive heat across Georgia, United States. *Weather Clim Soc*, *6*(2), 253-263. <https://doi.org/10.1175/wcas-d-13-00037.1>
- Mallen, E., Stone, B., & Lanza, K. (2019). A methodological assessment of extreme heat mortality modeling and heat vulnerability mapping in Dallas, Texas. *Urban Clim*, *30*, 100528. <https://doi.org/10.1016/j.uclim.2019.100528>
- McGeehin, M. A., & Mirabelli, M. (2001). The potential impacts of climate variability and change on temperature-related morbidity and mortality in the United States. *Environ Health Perspect*, *109*(Suppl 2), 185-189. <https://doi.org/10.1289/ehp.109-1240665>
- Medina-Ramón, M., & Schwartz, J. (2007). Temperature, temperature extremes, and mortality: a study of acclimatisation and effect modification in 50 US cities. *Occup Environ Med*, *64*(12), 827-833. <https://doi.org/10.1136/oem.2007.033175>
- Meier, H. C. S., & Mitchell, B. C. (2021). *Historic redlining scores for 2010 and 2020 US census tracts* Version 1) Inter-university Consortium for Political and Social Research. <https://doi.org/10.3886/E141121V1>
- Morello-Frosch, R., & Jesdale, B. M. (2006). Separate and unequal: residential segregation and estimated cancer risks associated with ambient air toxics in U.S. metropolitan areas. *Environ Health Perspect*, *114*(3), 386-393. <https://doi.org/10.1289/ehp.8500>
- Namin, S., Xu, W., Zhou, Y., & Beyer, K. (2020). The legacy of the Home Owners' Loan Corporation and the political ecology of urban trees and air pollution in the United States. *Soc Sci Med*, *246*, 112758. <https://doi.org/10.1016/j.socscimed.2019.112758>
- Nardone, A., Rudolph, K. E., Morello-Frosch, R., & Casey, J. A. (2021). Redlines and greenspace: the relationship between historical redlining and 2010 greenspace across the United States. *Environ Health Perspect*, *129*(1), 17006. <https://doi.org/10.1289/EHP7495>
- Naughton, M. P., Henderson, A., Mirabelli, M. C., Kaiser, R., Wilhelm, J. L., Kieszak, S. M., Rubin, C. H., & McGeehin, M. A. (2002). Heat-related mortality during a 1999 heat wave in Chicago. *Am J Prev Med*, *22*(4), 221-227. [https://doi.org/10.1016/s0749-3797\(02\)00421-x](https://doi.org/10.1016/s0749-3797(02)00421-x)
- Nawrot, T. S., Torfs, R., Fierens, F., De Henauw, S., Hoet, P. H., Van Kersschaever, G., De Backer, G., & Nemery, B. (2007). Stronger associations between daily mortality and fine particulate air pollution in summer than in winter: evidence from a heavily polluted

- region in western Europe [10.1136/jech.2005.044263]. *Journal of Epidemiology and Community Health*, 61(2), 146. <http://jech.bmj.com/content/61/2/146.abstract>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2465652/pdf/146.pdf>
- Nayak, S. G., Shrestha, S., Kinney, P. L., Ross, Z., Sheridan, S. C., Pantea, C. I., Hsu, W. H., Muscatiello, N., & Hwang, S. A. (2018). Development of a heat vulnerability index for New York State. *Public Health*, 161, 127-137.
<https://doi.org/10.1016/j.puhe.2017.09.006>
- Nelson, R. K., Winling, L., Marciano, R., Connolly, N., & Ayers, E. L. (2020). Mapping inequality: redlining in new deal America. *American panorama*.
<https://dsl.richmond.edu/panorama/redlining/#loc=5/39.1/-94.58&text=intro>
- O'Neill, M. S., Zanobetti, A., & Schwartz, J. (2003). Modifiers of the temperature and mortality association in seven US cities. *Am J Epidemiol*, 157(12), 1074-1082.
<https://doi.org/10.1093/aje/kwg096>
- O'Neill, M. S., Zanobetti, A., & Schwartz, J. (2005). Disparities by race in heat-related mortality in four US cities: the role of air conditioning prevalence. *J Urban Health*, 82(2), 191-197.
<https://doi.org/10.1093/jurban/jti043>
- Organization, W. H. (2021). *Climate change and health: vulnerability and adaptation assessment*. W. H. Organization.
<https://www.who.int/publications/i/item/9789240036383>
- Pearcy, M. (2020). "The most insidious legacy"—teaching about redlining and the impact of racial residential segregation. *Geography Teacher*, 17(2), 44-55.
<https://doi.org/10.1080/19338341.2020.1759118>
- Perry, R. J. (2007). *"Race" and Racism: The Development of Modern Racism in America*. Palgrave Macmillan.
- Petkova, E. P., Gasparini, A., & Kinney, P. L. (2014). Heat and mortality in New York City since the beginning of the 20th century. *Epidemiology*, 25(4), 554-560.
<https://doi.org/10.1097/EDE.000000000000123>
- Physical Housing Characteristics for Occupied Housing Units*. (2019). [CVS Data File]. U.S. Census Bureau.
<https://data.census.gov/cedsci/table?q=housing&g=0100000US%241400000&tid=ACSS T5Y2019.S2504>
- Poverty Status in the Past 12 Months*. (2019). [CVS Data File]. U.S. Census Bureau.
<https://data.census.gov/cedsci/table?q=poverty&g=0100000US%241400000&tid=ACSS T5Y2019.S1701>
- Prudent, N., Houghton, A., & Luber, G. (2016). Assessing climate change and health vulnerability at the local level: Travis County, Texas. *Disasters*, 40(4), 740-752.
<https://doi.org/10.1111/disa.12177>
- Qian, Z., He, Q., Lin, H.-M., Kong, L., Bentley, C. M., Liu, W., & Zhou, D. (2008). High temperatures enhanced acute mortality effects of ambient particle pollution in the "oven" city of Wuhan, China. *Environ Health Perspect*, 116(9), 1172.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2535618/pdf/ehp-116-1172.pdf>
- Reid, C. E., O'Neill, M. S., Gronlund, C. J., Brines, S. J., Brown, D. G., Diez-Roux, A. V., & Schwartz, J. (2009). Mapping community determinants of heat vulnerability. *Environ Health Perspect*, 117(11), 1730-1736. <https://doi.org/10.1289/ehp.0900683>
- Romanello, M., McGushin, A., Claudia Di, N., Drummond, P., Hughes, N., Jamart, L., Kennard, H., Lampard, P., Baltazar Solano, R., Arnell, N., Ayeb-Karlsson, S., Belesova, K., Cai,

- W., Campbell-Lendrum, D., Capstick, S., Chambers, J., Chu, L., Ciampi, L., Dalin, C., . . . Hamilton, I. (2021). The 2021 report of the Lancet Countdown on health and climate change: code red for a healthy future. *Lancet*, 398(10311), 1619-1662. [https://doi.org/10.1016/S0140-6736\(21\)01787-6](https://doi.org/10.1016/S0140-6736(21)01787-6)
- Schinasi, L. H., Kanungo, C., Christman, Z., Barber, S., Tabb, L., & Headen, I. (2022). Associations between historical redlining and present-day heat vulnerability housing and land cover characteristics in Philadelphia, PA. *J Urban Health*, 99, 134-145. <https://doi.org/10.1007/s11524-021-00602-6>
- Schwartz, J. (2005). Who is sensitive to extremes of temperature?: a case-only analysis. *Epidemiology*, 16(1), 67-72. <https://doi.org/10.1097/01.ede.0000147114.25957.71>
- Selected Economic Characteristics*. (2019). [CSV Data File]. U.S. Census Bureau. <https://data.census.gov/cedsci/table?q=economic&g=0100000US%241400000&tid=ACSDP5Y2019.DP03>
- Selected Social Characteristics in the United States*. (2019). [CSV Data File]. U.S. Census Bureau. <https://data.census.gov/cedsci/table?q=household&g=0100000US%241400000&tid=ACSDP5Y2019.DP02>
- Semenza, J. C., McCullough, J. E., Flanders, W. D., McGeehin, M. A., & Lumpkin, J. R. (1999). Excess hospital admissions during the July 1995 heat wave in Chicago. *Am J Prev Med*, 16(4), 269-277. [https://doi.org/10.1016/s0749-3797\(99\)00025-2](https://doi.org/10.1016/s0749-3797(99)00025-2)
- Semenza, J. C., Rubin, C. H., Falter, K. H., Selanikio, J. D., Flanders, W. D., Howe, H. L., & Wilhelm, J. L. (1996). Heat-related deaths during the July 1995 heat wave in Chicago. *N Engl J Med*, 335(2), 84-90. <https://doi.org/10.1056/NEJM199607113350203>
- Shen, H., Li, X., Cheng, Q., Zeng, C., Yang, G., Li, H., & Zhang, L. (2015). Missing information reconstruction of remote Sensing data: a technical review. *IEEE Trans Geosci Remote Sens*, 3(3), 61-85. <https://doi.org/10.1109/MGRS.2015.2441912>
- Smoyer, K. E. (1998). Putting risk in its place: methodological considerations for investigating extreme event health risk. *Soc Sci Med*, 47(11), 1809-1824. [https://doi.org/10.1016/S0277-9536\(98\)00237-8](https://doi.org/10.1016/S0277-9536(98)00237-8)
- Spangler, K. R., & Wellenius, G. A. (2020). Spatial patterns of recent US summertime heat trends: Implications for heat sensitivity and health adaptations. *Environ Res Commun*, 2(3), 035002. <https://doi.org/10.1088/2515-7620/ab7abb>
- Thornton, M. M., Shrestha, R., Wei, Y., Thornton, P. E., Kao, S., & Wilson, B. E. (2020). *Daymet: Daily Surface Weather Data on a 1-km Grid for North America, Version 4*. <https://doi.org/10.3886/E110663V1>
- Uejio, C. K., Wilhelmi, O. V., Golden, J. S., Mills, D. M., Gulino, S. P., & Samenow, J. P. (2011). Intra-urban societal vulnerability to extreme heat: the role of heat exposure and the built environment, socioeconomics, and neighborhood stability. *Health Place*, 17(2), 498-507. <https://doi.org/10.1016/j.healthplace.2010.12.005>
- Whitman, S., Good, G., Donoghue, E. R., Benbow, N., Shou, W., & Mou, S. (1997). Mortality in Chicago attributed to the July 1995 heat wave. *Am J Public Health*, 87(9), 1515-1518. <https://doi.org/10.2105/ajph.87.9.1515>
- Wilson, B. (2020). Urban heat management and the legacy of redlining. *J Am Plann Assoc*, 86(4), 443-457. <https://doi.org/10.1080/01944363.2020.1759127>
- Wold, S., Esbensen, K., & Geladi, P. (1987). Principal component analysis. *Chemometrics Intelligent Lab Syst*, 2(1), 37-52. [https://doi.org/10.1016/0169-7439\(87\)80084-9](https://doi.org/10.1016/0169-7439(87)80084-9)

- Wolf, T., & McGregor, G. (2013). The development of a heat wave vulnerability index for London, United Kingdom. *Weather Clim Extremes*, *1*, 59-68.
<https://doi.org/10.1016/j.wace.2013.07.004>
- Xu, Y., Dadvand, P., Barrera-Gómez, J., Sartini, C., Marí-Dell'Olmo, M., Borrell, C., Medina-Ramón, M., Sunyer, J., & Basagaña, X. (2013). Differences on the effect of heat waves on mortality by sociodemographic and urban landscape characteristics. *J Epidemiol Community Health*, *67*(6), 519-525. <https://doi.org/10.1136/jech-2012-201899>
- Xu, Z., Fitzgerald, G., Guo, Y., Jalaludin, B., & Tong, S. (2016). Impact of heatwave on mortality under different heatwave definitions: A systematic review and meta-analysis. *Environ Int*, *89-90*, 193-203. <https://doi.org/10.1016/j.envint.2016.02.007>
- Yin, Q., Wang, J., Ren, Z., Li, J., & Guo, Y. (2019). Mapping the increased minimum mortality temperatures in the context of global climate change. *Nat Commun*, *10*(1), 4640.
<https://doi.org/10.1038/s41467-019-12663-y>
- Zanobetti, A., O'Neill, M. S., Gronlund, C. J., & Schwartz, J. D. (2013). Susceptibility to mortality in weather extremes: effect modification by personal and small-area characteristics. *Epidemiology*, *24*(6), 809-819.
<https://doi.org/10.1097/01.ede.0000434432.06765.91>
- Zanobetti, A., & Schwartz, J. (2008). Temperature and mortality in nine US cities. *Epidemiology*, *19*(4), 563-570. <https://doi.org/10.1097/EDE.0b013e31816d652d>
- Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., & Huang, M. (2021). IPCC 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. In: Cambridge University Press.
- Zou, Y., Wang, Y., Zhang, Y., & Koo, J.-H. (2017). Arctic sea ice, Eurasia snow, and extreme winter haze in China [10.1126/sciadv.1602751]. *Science Advances*, *3*(3).
<http://advances.sciencemag.org/content/3/3/e1602751.abstract>